

DERIVATIVES OF MEROMORPHIC FUNCTIONS WITH MULTIPLE ZEROS AND ELLIPTIC FUNCTIONS

PAI YANG, SHAHAR NEVO, AND XUECHENG PANG

ABSTRACT. Let f be a nonconstant meromorphic function in the plane and h be a nonconstant elliptic function. We show that if all zeros of f are multiple except finitely many and $T(r, h) = o\{T(r, f)\}$ as $r \rightarrow \infty$, then $f' = h$ has infinitely many solutions (including poles).

1. INTRODUCTION

Recall that an elliptic function [1] is a meromorphic function h defined on $\mathbb{C}$ for which there exist two non-zero complex numbers ω_1 and ω_2 with ω_1/ω_2 not real such that $h(z + \omega_1) = h(z + \omega_2) = h(z)$ for all z on $\mathbb{C}$.

We use the following notation. For $z_0 \in \mathbb{C}$ and $r > 0$, $\Delta(z_0, r) = \{z : |z - z_0| < r\}$, $\Delta'(z_0, r) = \{z : 0 < |z - z_0| < r\}$, $\Delta = \Delta(0, 1)$, $\overline{\Delta}(0, r) = \{z : |z| \leq r\}$, and $\Gamma(0, r) = \{z : |z| = r\}$. For f meromorphic in a domain D , we denote

$$S(D, f) = \frac{1}{\pi} \iint_D [f^\#(z)]^2 dx dy, \quad S(r, f) = S(\Delta(0, r), f).$$

Here

$$f^\#(z) = \frac{|f'(z)|}{1 + |f(z)|^2}$$

denotes the spherical derivative. We use the Ahlfors-Shimizu form of the Nevanlinna characteristic function, given by

$$T_0(r, f) = \int_0^r \frac{S(t, f)}{t} dt.$$

Let $n(r, f)$ denote the number of poles of $f(z)$ in $\Delta(0, r)$ (counting multiplicity). We write $f_n \xrightarrow{x} f$ in D to indicate that the sequence $\{f_n\}$ converges to f in the spherical metric uniformly on compact subsets of D and $f_n \Rightarrow f$ in D if the convergence is in the Euclidean metric.

In 1959, Hayman [7] proved the following seminal result, which has come to be known as Hayman's Alternative.

Theorem A. *Let f be a transcendental meromorphic function on the complex plane $\mathbb{C}$. Then either*

- (i) *f assumes each value $a \in \mathbb{C}$ infinitely often, or*

2010 *Mathematics Subject Classification.* 30D30, 30D35, 30D45.

Key words and phrases. Normal family, elliptic function.

(ii) $f^{(k)}$ assumes each value $b \in \mathbb{C} \setminus \{0\}$ infinitely often for $k = 1, 2, \dots$

Obviously, if f is a transcendental meromorphic function such that $f \neq 0$ for all $z \in \mathbb{C}$, then $f^{(k)}$ assumes each value $b \in \mathbb{C} \setminus \{0\}$ infinitely often for $k = 1, 2, \dots$. In recent years, it has become clear that, in many instances, the condition $f \neq 0$ can be replaced by the assumption that all zeros of f have sufficiently high multiplicity. This announcement concerns such extension of Theorem A. We restrict our attention to the case $k = 1$.

Before stating our result, let us present several theorems that show results already obtained in this subject.

Theorem B. [23, Theorem 3] *Let f be a transcendental meromorphic function on $\mathbb{C}$, all of whose zeros have multiplicity at least 3. Then f' assumes each nonzero complex value infinitely often.*

The analogue of Theorem B with 3 replaced by 2 does hold for functions of finite order.

Theorem C. [23, Lemma 6] *Let f be a transcendental meromorphic function of finite order on $\mathbb{C}$, all of whose zeros are multiple. Then f' assumes every finite non-zero value infinitely often.*

This is an instant corollary of Theorem A and the following important result.

Theorem D. [3, Theorem 3] *Let f be a transcendental meromorphic function of finite order on $\mathbb{C}$ with an infinite number of multiple zeros. Then f' assumes every finite non-zero value infinitely often.*

Indeed, if in Theorem C, f vanishes only finitely often, then f' must take on every nonzero value infinitely often by Theorem A; Otherwise, Theorem D implies the same conclusion.

Bergweiler and Eremenko gave a counter example in [3] which shows Theorem D is not true in general for functions of infinite order.

In 2006, Shahar Nevo, Xuecheng Pang and Lawrence Zalcman gave the following extension of Theorem B and Theorem C.

Theorem E. [19, Theorem 1] *Let f be a transcendental meromorphic function on $\mathbb{C}$, all but finitely many of whose zeros are multiple, and let $R \neq 0$ be a rational function. Then $f' - R$ has infinitely many zeros.*

In this paper, we continue to study the value distribution of the derivative of meromorphic functions with multiple zeros. We have the following theorem.

Theorem 1. *Let f be a nonconstant meromorphic function on $\mathbb{C}$ and h be a non-constant elliptic function. Then if all zeros of f are multiple except finitely many and $T_0(r, h) = o\{T_0(r, f)\}$ as $r \rightarrow \infty$, then $f' = h$ has infinitely many solutions (including the possibility of infinitely many common poles of f and h).*

Remark. $T(r, f)$ denotes the usual Nevanlinna characteristic function. Since $T(r, f) - T_0(r, f)$ is bounded as a function of r , one can replace $T_0(r, f)$ with $T(r, f)$ in Theorem 1.

2. PRELIMINARY RESULTS

Lemma 2.1. [21, Theorem 1][22, Theorem VI.21] *Let f be a meromorphic function in Δ , and let a_1, a_2, a_3 be 3 distinct complex numbers. Assume that the number of zeros of $\prod_{i=1}^3 (f(z) - a_i)$ in Δ is $\leq n$, where multiple zeros are counted only once, then*

$$S(r, f) \leq n + \frac{A}{1-r}, \quad 0 \leq r < 1,$$

and $A > 0$ is a constant, which depends on a_1, a_2, a_3 only.

Lemma 2.2. *If $h(z)$ is a nonconstant elliptic function with primitive periods ω_1, ω_2 , where ω_1/ω_2 not real, then*

$$T_0(r, h) = Ar^2(1 + o(1)) \quad \text{as } r \rightarrow \infty,$$

where $A > 0$ is a constant.

This follows from [2, Corollary 2].

Lemma 2.3. [20, Theorem 3][11, Theorem 1] *Let $\psi \neq 0$ be a meromorphic function in a domain $D \subset \mathbb{C}$ and $k \in \mathbb{N}$. Let $\mathcal{F}$ be a family of meromorphic functions in D , such that f and $f^{(k)} - \psi$ have no zeros and f and ψ have no common poles for each $f \in \mathcal{F}$. Then $\mathcal{F}$ is normal on D .*

Lemma 2.4. *Let f be a nonconstant meromorphic function of finite order on $\mathbb{C}$, all of whose zeros are multiple. If $f'(z) \neq 1$ on $\mathbb{C}$, then*

$$f(z) = \frac{(z-a)^2}{z-b}$$

for some a and $b (\neq a)$ on $\mathbb{C}$.

This follows from Lemma 6 (with $j = 1$ and $k = 2$) and Lemma 8 (with $k = 1$) of [23].

Lemma 2.5. [15, Lemma 2] *Let $\mathcal{F}$ be a family of functions meromorphic in a domain D , all of whose zeros have multiplicity at least k , and suppose that there exists $A \geq 1$ such that $|f^{(k)}(z)| \leq A$ whenever $f(z) = 0$. Then if $\mathcal{F}$ is not normal at z_0 , there exist, for each $0 \leq \alpha \leq k$,*

- (a) *points $z_n, z_n \rightarrow z_0$;*
- (b) *functions $f_n \in \mathcal{F}$; and*
- (c) *positive numbers $\rho_n \rightarrow 0$*

such that $\rho_n^{-\alpha} f_n(z_n + \rho_n \zeta) = g_n(\zeta) \xrightarrow{x} g(\zeta)$ on $\mathbb{C}$, where g is a nonconstant meromorphic function on $\mathbb{C}$, all of whose zeros have multiplicity at least k , such that $g^\#(\zeta) \leq g^\#(0) = kA + 1$.

Lemma 2.6. [16, Lemma 3.1] *Let $\{f_n\}$ be a family of functions meromorphic in $\Delta(z_0, r)$, all of whose zeros and poles are multiple; and let $\{b_n\}$ be a sequence of holomorphic functions in $\Delta(z_0, r)$ such that $b_n \Rightarrow b$ in $\Delta(z_0, r)$, where $b(z) \neq 0, z \in \Delta(z_0, r)$. If $f'_n(z) \neq b_n(z)$ for $z \in \Delta(z_0, r)$, then $\{f_n\}$ is normal in $\Delta(z_0, r)$.*

Lemma 2.7. [17, Lemma 3.1] *Let $\{f_n\}$ be a sequence of meromorphic functions in $\Delta(z_0, r)$ and $\{\psi_n\}$ a sequence of holomorphic functions in $\Delta(z_0, r)$ such that $\psi_n \Rightarrow \psi$, where $\psi(z) \neq 0$ in $\Delta(z_0, r)$. If for each n , $f_n(z) \neq 0$ and $f'_n(z) \neq \psi_n(z)$ for $z \in \Delta(z_0, r)$, then $\{f_n\}$ is normal in $\Delta(z_0, r)$.*

Lemma 2.8. [17, Lemma 3.2] *Let E be a (countable) discrete set in $\Delta(z_0, r)$ which has no accumulation points in $\Delta(z_0, r)$ and let $\{\psi_n\}$ be a sequence of holomorphic functions in $\Delta(z_0, r)$ such that $\psi_n \Rightarrow \psi$ in $\Delta(z_0, r)$, where $\psi \neq 0, \infty$ in $\Delta(z_0, r)$. Let $\{f_n\}$ be a sequence of functions meromorphic in $\Delta(z_0, r)$, all of whose zeros are multiple, such that $f'_n(z) \neq \psi_n(z)$ for all n and all $z \in \Delta(z_0, r)$. Let $a_1 \in E$ and suppose that*

- (a) *for some $a_1 \in E$, no subsequence of $\{f_n\}$ is normal at a_1 ;*
- (b) *there exists $\delta > 0$ such that each f_n has a single (multiple) zero in $\Delta(a_1, \delta)$; and*
- (c) *$f_n(z) \xrightarrow{X} f(z)$ in $\Delta(z_0, r) \setminus E$.*

Then

- (d) *there exists $\eta_0 > 0$ such that for each $0 < \eta < \eta_0$, f_n has a single simple pole in $\Delta(a_1, \eta)$ for sufficiently large n ; and*
- (e) *$f(z) = \int_{a_1}^z \psi(\zeta) d\zeta$.*

Lemma 2.9. [17, Lemma 3.7] *Let $\{f_n\}$ be a sequence of functions meromorphic on $\Delta(z_0, r)$, all of whose zeros are multiple, and let ψ be a non-vanishing holomorphic function in $\Delta(z_0, r)$. Suppose that*

- (a) *$\{f_n\}$ is quasinormal in $\Delta(z_0, r)$;*
- (b) *$f'_n(z) \neq \psi(z)$ for $z \in \Delta(z_0, r)$ and $n = 1, 2, 3, \dots$;*
- (c) *no subsequence of $\{f_n(z)\}$ is normal at 0.*

Then there exists $0 < \delta < r$ such that f_n has only a single (multiple) zero in $\Delta(0, \delta)$ for sufficiently large n .

Lemma 2.10. *Let $\{f_n\}$ be a family of meromorphic functions on the plane domain D , all whose zeros are multiple, If there exists a holomorphic function φ univalent in D such that $f'_n(z) \neq \varphi'(z)$ for each $n \in \mathbb{N}$ and $z \in D$, Then $\{f_n\}$ is quasinormal of order 1 in D .*

This follows from Theorem 1 (with $k = 1$ and $\mathcal{F} = \{f_n\}$) of [14].

Lemma 2.11. [5, Lemma 12] *Let R be a nonconstant rational function satisfying $R' \neq 0$ on $\mathbb{C}$. Then $R(z) = az + b$ or $R(z) = \frac{a}{(z+c)^n} + b$, where $n \in \mathbb{N}$ and $a \neq 0, b, c \in \mathbb{C}$.*

3. AUXILIARY LEMMAS

Lemma 3.1. *Let $k \geq 2$ be an integer and let R be a rational function on $\mathbb{C}$. Suppose that $R'(z) \neq \frac{1}{z^k}$. Then R is a constant function.*

Proof. Since $R'(z) \neq \frac{1}{z^k}$, we have $R(0) \neq \infty$ and hence

$$R'(z) - \frac{1}{z^k} = \left(R(z) - \frac{1}{1-k} \frac{1}{z^{k-1}} \right)' \neq 0.$$

By Lemma 2.11, either $R(z) = az + b + \frac{1}{1-k} \frac{1}{z^{k-1}}$, or $R(z) = \frac{a}{(z+c)^n} + b + \frac{1}{1-k} \frac{1}{z^{k-1}}$. If $R(z) = az + b + \frac{1}{1-k} \frac{1}{z^{k-1}}$, then we have $R(0) = \infty$. A contradiction. Thus, $R(z) = \frac{a}{(z+c)^n} + b + \frac{1}{1-k} \frac{1}{z^{k-1}}$. Since $R(0) \neq \infty$, we have that $c = 0$, $n = k - 1$, $a = \frac{1}{k-1}$ and $R(z) \equiv b$. $\square$

Lemma 3.2. *Let k be a positive integer and let R be a rational function satisfying $R'(z) \neq z^k$ on $\mathbb{C}$. If all zeros of R are multiple, then*

$$(3.1) \quad R(z) = \frac{\prod_{i=1}^{n+k+1} (z - \alpha_i)}{(k+1)(z - \beta)^n},$$

where n is a nonnegative integer $\beta \in \mathbb{C}$ and $\alpha_i (\neq 0, \beta)$, for $1 \leq i \leq n + k + 1$ and every zero α_i is counted due to multiplicity.

Proof. Obviously, $(R(z) - \frac{z^{k+1}}{k+1})' \neq 0$ and since all the zeros of R are multiple, then $R(z) - \frac{z^{k+1}}{k+1}$ is a nonconstant rational function. By Lemma 2.11, $R(z) = \frac{z^{k+1}}{k+1} + az + b$ or $R(z) = \frac{z^{k+1}}{k+1} + \frac{a}{(z+c)^n} + b$, where $n \in \mathbb{N}$ and $a \neq 0, b, c \in \mathbb{C}$. Let us consider separately these two cases.

Case 1: $R(z) = \frac{z^{k+1}}{k+1} + az + b$.

Let z_0 be a zero of $R(z)$, then $R(z_0) = 0$ and hence $R'(z_0) = 0$, i.e.,

$$(3.2) \quad \begin{aligned} R(z_0) &= \frac{1}{k+1} z_0^{k+1} + az_0 + b = 0, \\ R'(z_0) &= z_0^k + a = 0. \end{aligned}$$

It follows that $z_0 = -\frac{b(k+1)}{ak}$ and hence

$$(3.3) \quad R(z) = \frac{1}{k+1} \left(z + \frac{b(k+1)}{ak} \right)^{k+1}.$$

Comparing the coefficients of (3.2) and (3.3), we have $k = 1$ and hence $R(z) = \frac{(z-\alpha)^2}{2}$, where $\alpha = -\frac{2b}{a}$. Now $R(z)$ have the form of (3.1) with $\alpha_1 = \alpha_2 = \alpha$, $k = 1$ and $n = 0$.

Case 2: $R(z) = \frac{1}{k+1} z^{k+1} + \frac{a}{(z+c)^n} + b$.

Let $\{\alpha_1, \alpha_2, \dots, \alpha_{n+k+1}\}$ be the set of the zeros of $R(z)$ and $\beta = -c$. Obviously, $R(z)$ have the form of (3.1).

Since $R'(z) \neq z^k$, $R'(0) \neq 0$, and since all zeros of R are multiple, $R(0) \neq 0$. Hence $\alpha_i \neq 0$ for $1 \leq i \leq n + k + 1$. $\square$

Lemma 3.3. *Let $\{f_n(z)\}$ be a family of meromorphic functions in Δ , all of whose zeros are multiple. Let $F_n(z) = z^k f_n(z)$, where $k \neq 0$ is an integer. Let $\{b_n(z)\}$ be a sequence of holomorphic functions in Δ such that $b_n(z) \Rightarrow b(z)$ in Δ , where $b(z) \neq 0$ is a holomorphic function in Δ . Suppose that for each n , $f'_n(z) \neq z^{-k} b_n(z)$ and $\{f_n(z)\}$ is not normal at 0 and normal in Δ' . Then $\{F_n(z)\}$ is normal in Δ' but not normal at 0.*

Proof. It is obvious that $\{F_n(z)\}$ is normal in Δ' . Suppose that $\{F_n(z)\}$ is normal at 0.

Case 1: k is a positive integer.

Since $f'_n(z) \neq z^{-k}b_n(z)$ and $b_n(z) \Rightarrow b(z)$ on Δ , and $b(0) \neq 0$ we have, for sufficiently large n , $f'_n(0) \neq \infty$ and $f_n(0) \neq \infty$. Obviously, we have $F_n(0) = 0$. Without loss of generality, we can suppose that for all n , $f_n(0) \neq \infty$ and $F_n(0) = 0$. Since $\{F_n(z)\}$ is normal at 0 and $F_n(0) = 0$, there exists $\delta > 0$ such that for all n , $|F_n(z)| \leq 1$ in $\Delta(0, \delta)$. Thus $f_n(0) \neq \infty$ in $\Delta'(0, \delta)$, and hence f_n is holomorphic in $\Delta(0, \delta)$. Now, we have

$$|f_n(z)| = \frac{|F_n(z)|}{|z|^k} \leq \left(\frac{1}{\delta}\right)^k, \quad z \in \Gamma(0, \delta).$$

By the maximum principle, this holds throughout $\Delta(0, \delta)$. It follows that $\{f_n\}$ is normal. A contradiction.

Case 2: k is a negative integer.

Let $m = -k$. Since $f'_n(z) \neq z^m b_n(z)$ and $b_n(z) \Rightarrow b(z)$ in Δ , we have, for sufficiently large n , $f'_n(0) \neq 0$ and hence $f_n(0) \neq 0$. Obviously, we have $F_n(0) = \infty$. Without loss of generality, we can suppose that for all n , $f_n(0) \neq 0$ and $F_n(0) = \infty$. Since $\{F_n(z)\}$ is normal at 0 and $F_n(0) = \infty$, there exists $\delta > 0$ such that for all n , $|F_n(z)| \geq 1$ on $\Delta(0, \delta)$. Thus $f_n(z) \neq 0$ in $\Delta'(0, \delta)$, and hence $\frac{1}{f_n}$ is holomorphic in $\Delta(0, \delta)$. Now, we have

$$\left| \frac{1}{f_n(z)} \right| = \left| \frac{1}{F_n(z)} \frac{1}{z^m} \right| \leq \frac{1}{\delta^m}, \quad z \in \Gamma(0, \delta).$$

By the maximum principle, this holds throughout $\Delta(0, \delta)$. It follows that $\{f_n\}$ is normal. A contradiction. $\square$

Lemma 3.4. *Let $\{f_n\}$ and $\{\psi_n\}$ be two sequences of functions meromorphic on the plane domain D . Let $f(z)$ and $\psi(z)$ be two meromorphic functions in D . Suppose that*

- (a) $f_n(z) \xrightarrow{X} f(z)$ in D and $\psi_n(z) \xrightarrow{X} \psi(z)$ in D ;
- (b) $f'_n(z) \neq \psi_n(z)$ in D .

Then, either $f'(z) \equiv \psi(z)$ in D , or $f'(z) \neq \psi(z)$ in D .

Proof. Suppose that there exists $z_0 \in D$ such that $f'(z_0) = \psi(z_0)$.

Let us separate into two cases.

Case 1: $\psi(z_0) \neq \infty$.

There exists $r > 0$ such that f, ψ, f_n, ψ_n are analytic in $\Delta(z_0, r)$ for large enough n . We have that $f'_n - \psi_n \Rightarrow f' - \psi$ in $\Delta(z_0, r)$, and by condition (b) and Hurwitz's Theorem $f' - \psi \equiv 0$ in $\Delta(z_0, r)$ and so in all of D . In this case, the first possibility occurs.

Case 2: $\psi(z_0) = \infty$.

Suppose by negation that $f' - \psi$ is not constantly zero. Thus, for small enough r , $\frac{1}{f' - \psi}$ is analytic in $\Delta'(z_0, r)$, and since $\frac{1}{f'_n - \psi_n}$ are analytic in D and $\frac{1}{f'_n - \psi_n} \Rightarrow \frac{1}{f' - \psi}$ in $\Delta'(z_0, r)$, we get by the maximum principle that $\left\{ \frac{1}{f'_n - \psi_n} \right\}_{n=1}^{\infty}$ is a uniformly convergent

Cauchy sequence of analytic functions in $\Delta(z_0, r)$ which thus converges (uniformly) to an analytic function there. Thus $\frac{1}{f' - \psi}$ is extended analytically to $\Delta(z_0, r)$. Let $m_1, m_2 \geq 1$ be the orders of the pole of f and ψ at z_0 , respectively. A pole of $f' - \psi$ at z_0 , if it exists, is of order at most $\max\{m_1 + 1, m_2\}$. This is also the maximal multiplicity of zero of $\frac{1}{f' - \psi}$ at z_0 . (Observe that it is possible that $f'(z_0) - \psi(z_0)$ is a finite value, but this value cannot be 0 since $\frac{1}{f' - \psi}$ is analytic at z_0). Now, for every $0 < \delta < r$, f_n has, for large enough n , m_1 poles (counting multiplicities) in $\Delta(z_0, \delta)$, and ψ_n has m_2 poles in $\Delta(z_0, \delta)$ (counting multiplicities). Since f_n and ψ_n have no common poles, then $\frac{1}{f'_n - \psi_n}$ has at least $m_1 + m_2 + 1$ zeros in $\Delta(z_0, \delta)$. But $m_1 + m_2 + 1 \geq \max\{m_1 + 1, m_2\}$ and this is a contradiction. Thus $f' - \psi \equiv 0$. So also in the case of a common pole, we have that $f' \equiv \psi$ in D and the lemma is proved. $\square$

Lemma 3.5. *Let $\{f_n\}$ be a family of meromorphic functions in Δ , all of whose zeros are multiple. Let $\{b_n\}$ be a sequence of holomorphic functions in Δ such that $b_n \Rightarrow 1$ in Δ , and let k be a positive integer. Suppose that*

- (a) $f'_n(z) \neq z^k b_n(z)$ in Δ ;
- (b) *there exists points $z_n \rightarrow 0$ such that $f_n(z_n) = 0$;*
- (c) $f_n(z) \xrightarrow{X} f(z)$ in Δ' , where $f(z)$ is a meromorphic function in Δ' .

Then $f(z) = \frac{z^{k+1+c}}{k+1}$ in Δ' , where c is a constant.

Proof. Let $F_n(z) = \frac{f'_n(z)}{z^k}$. Since $b_n \Rightarrow 1$ in Δ and (a), we have for sufficiently large n , $f'_n(0) \neq 0$ and hence $F_n(0) = \infty$. Without loss of generality, we may assume that for all n , $f'_n(0) \neq 0$ and $F_n(0) = \infty$. Since all zeros of $\{f_n(z)\}$ are multiple, $f_n(0) \neq 0$. Hence we have $z_n \neq 0$ and $F_n(z_n) = 0$.

We claim that $\{F_n(z)\}$ is not normal at 0 and hence $\{\frac{f'_n(z)}{z^k b_n(z)} - 1\}$ is also not normal at 0. Indeed, since $F_n(z_n) = 0$ and $F_n(0) = \infty$, the family $\{F_n(\zeta)\}$ is not equicontinuous at 0 and hence cannot be normal at 0.

By (a) and (c), we have

$$0 \neq \frac{f'_n(z)}{z^k b_n(z)} - 1 \Rightarrow \frac{f'(z)}{z^k} - 1, \quad z \in \Delta' - \{f^{-1}(\infty)\}.$$

By Hurwitz's Theorem, either $\frac{f'(z)}{z^k} - 1 \equiv 0$ in Δ' , or $\frac{f'(z)}{z^k} - 1 \neq 0$ in Δ' . Suppose that $\frac{f'(z)}{z^k} - 1 \neq 0$ in Δ' . Since $f(z)$ is a meromorphic function, then there exists $\delta > 0$ such that $f(z)$ has no poles on $\Gamma(0, \delta)$ and $f'_n(z)$ converges uniformly to $f'(\zeta)$ on $\Gamma(0, \delta)$. Now, we have

$$\infty \neq \frac{1}{\frac{f'_n(z)}{z^k b_n(z)} - 1} \Rightarrow \frac{1}{\frac{f'(z)}{z^k} - 1}, \quad z \in \Gamma(0, \delta).$$

As in the proof of Lemma 3.4, since the function in the left hand side is holomorphic, we have by the maximum principle that this holds throughout $\Delta(0, \delta)$. So $\{\frac{f'_n(z)}{z^k b_n(z)} - 1\}$ is normal at 0. A contradiction. Thus, $\frac{f'(z)}{z^k} - 1 \equiv 0$ in Δ' . Then $f(z) = \frac{z^{k+1+c}}{k+1}$ in Δ' , where c is a constant. $\square$

Lemma 3.6. *Let $\{f_n\}$ be a sequence of functions meromorphic in $\Delta(z_0, r)$. Suppose there exists $M_1 > 0$ such that for each n , $n(r, \frac{1}{f_n}) < M_1$. Suppose that $f_n \xrightarrow{x} f$ in $\Delta'(z_0, r)$, where f is a nonconstant meromorphic function or $f \equiv \infty$ in $\Delta'(z_0, r)$. Then there exists $M_2 > 0$ such that, for sufficiently large n ,*

$$S(\frac{r}{2}, f_n) < M_2.$$

Proof. Without loss of generality, we assume that $r = 1$ and so $\Delta(z_0, r) = \Delta$. Obviously, $\frac{1}{f_n} - 1 \xrightarrow{x} \frac{1}{f} - 1$ in Δ' and $\frac{1}{f} - 1 \not\equiv 0, \infty$ in Δ' . Then there exists $\frac{1}{2} < r < 1$ such that $\frac{1}{f} - 1$ has no poles and zeros in $\Gamma(0, r)$. Obviously, for sufficiently large n ,

$$\begin{aligned} n\left(r, \frac{1}{f_n - 1}\right) - n\left(r, \frac{1}{f_n}\right) &= n\left(r, \frac{1}{\frac{1}{f_n} - 1}\right) - n\left(r, \frac{1}{f_n} - 1\right) \\ &= \frac{1}{2\pi i} \int_{\Gamma(0, r)} \frac{(\frac{1}{f_n} - 1)'}{\frac{1}{f_n} - 1} dz \rightarrow \frac{1}{2\pi i} \int_{\Gamma(0, r)} \frac{(\frac{1}{f} - 1)'}{\frac{1}{f} - 1} dz. \end{aligned}$$

Set $M_3 = \frac{1}{2\pi i} \int_{\Gamma(0, r)} \frac{(\frac{1}{f} - 1)'}{\frac{1}{f} - 1} dz + n(r, \frac{1}{f_n}) + 1$. For sufficiently large n ,

$$\begin{aligned} \left(\frac{1}{2}, \frac{1}{f_n - 1}\right) &\leq n\left(r, \frac{1}{f_n - 1}\right) = n\left(r, \frac{1}{f_n}\right) + \int_{\Gamma(0, r)} \frac{(\frac{1}{f} - 1)'}{\frac{1}{f} - 1} dz + \varepsilon_n \\ &< M_1 + \int_{\Gamma(0, r)} \frac{(\frac{1}{f} - 1)'}{\frac{1}{f} - 1} dz = M_3 - 1 < M_3. \end{aligned}$$

(Here $\varepsilon_n \rightarrow 0$, but since the other terms are integers, then $\varepsilon_n = 0$ for large enough n .)

Obviously, $\frac{1}{f_n} - \frac{1}{2} \xrightarrow{x} \frac{1}{f} - \frac{1}{2}$ in Δ' and $\frac{1}{f} - \frac{1}{2} \not\equiv 0, \infty$ in Δ' . Then there exists $\frac{1}{2} < t < 1$ such that $\frac{1}{f} - \frac{1}{2}$ has no poles and zeros in $\Gamma(0, t)$. Clearly, for sufficiently large n ,

$$\begin{aligned} n\left(t, \frac{1}{f_n - 2}\right) - n\left(t, \frac{1}{f_n}\right) &= n\left(t, \frac{1}{\frac{1}{f_n} - \frac{1}{2}}\right) - n\left(t, \frac{1}{f_n} - \frac{1}{2}\right) \\ &= \frac{1}{2\pi i} \int_{\Gamma(0, t)} \frac{(\frac{1}{f_n} - \frac{1}{2})'}{\frac{1}{f_n} - \frac{1}{2}} dz \rightarrow \frac{1}{2\pi i} \int_{\Gamma(0, t)} \frac{(\frac{1}{f} - \frac{1}{2})'}{\frac{1}{f} - \frac{1}{2}} dz. \end{aligned}$$

Set $M_4 = \frac{1}{2\pi i} \int_{\Gamma(0, t)} \frac{(\frac{1}{f} - \frac{1}{2})'}{\frac{1}{f} - \frac{1}{2}} dz + n(t, \frac{1}{f_n}) + 1$. Similarly to the previous paragraph, we have, for sufficiently large n ,

$$n\left(\frac{1}{2}, \frac{1}{f_n - 2}\right) \leq n\left(t, \frac{1}{f_n - 2}\right) < M_4 - 1.$$

Let $M_2 = M_1 + M_3 + M_4 + 2A$. By Lemma 2.1, for sufficiently large n ,

$$S\left(\frac{1}{2}, f_n\right) \leq n\left(\frac{1}{2}, \frac{1}{f_n}\right) + n\left(\frac{1}{2}, \frac{1}{f_n - 1}\right) + n\left(\frac{1}{2}, \frac{1}{f_n - 2}\right) + 2A < M_2.$$

□

Lemma 3.7. *Let $\Psi(z)$ be a holomorphic univalent function on $\Delta(0, R)$ and $\{\Psi_n(z)\}$ be a sequence of holomorphic functions in $\Delta(0, R)$ such that $\Psi_n(z) \Rightarrow \Psi(z)$ in $\Delta(0, R)$. Then for each $r \in (0, R)$, we have, for sufficiently large n ,*

- (a) $\Psi_n(z)$ is a holomorphic univalent function in $\Delta(0, r)$;
- (b) there exists $\delta_1 > 0$ such that $\Delta(\Psi(0), \delta_1) \subset \Psi(\Delta(0, r))$ and $\Delta(\Psi(0), \delta_1) \subset \Psi_n(\Delta(0, r))$;
- (c) there exists $\delta_2 > 0$ such that $\Psi(\Delta(0, \delta_2)) \subset \Delta(\Psi(0), \delta_1)$ and $\Psi_n(\Delta(0, \delta_2)) \subset \Delta(\Psi(0), \delta_1)$.

Proof. Let $r < r_1 < R$. Suppose that there exists a subsequence of $\{\Psi_n(z)\}$ (that we continue to call $\{\Psi_n(z)\}$) which is not univalent in $\Delta(0, r)$. Then there exist distinct complex numbers $z_{n,1}$ and $z_{n,2}$ in $\Delta(0, r)$ such that $\Psi_n(z_{n,1}) = \Psi_n(z_{n,2})$. Obviously, $n(r_1, \frac{1}{\Psi_n(z) - \Psi_n(z_{n,1})}) \geq 2$. Without loss of generality, we may suppose that $z_{n,1} \rightarrow z_1$ and $z_{n,2} \rightarrow z_2$ as $n \rightarrow \infty$. Obviously $|z_1| \leq r, |z_2| \leq r$. Since $\Psi_n(z) \Rightarrow \Psi(z)$ in $\Delta(0, R)$, we have

$$\Psi(z_1) = \lim_{n \rightarrow \infty} \Psi_n(z_{n,1}) = \lim_{n \rightarrow \infty} \Psi_n(z_{n,2}) = \Psi(z_2).$$

Hence we have $z_1 = z_2$.

Obviously, $\Psi_n(z) - \Psi_n(z_{n,1}) \Rightarrow \Psi(z) - \Psi(z_1)$ in $\Delta(0, R)$ and $\Psi(z) - \Psi(z_1) \neq 0$ on $\Gamma(0, r_1)$. By the argument principle, for sufficiently large n ,

$$n \left(r_1, \frac{1}{\Psi_n(z) - \Psi_n(z_{n,1})} \right) = n \left(r_1, \frac{1}{\Psi(z) - \Psi(z_1)} \right) = 1.$$

A contradiction occurs and hence (a) holds.

Obviously, there exists δ_1 such that $\Delta(\Psi(0), 2\delta_1) \subset \Psi(\Delta(0, r))$, and hence $\Delta(\Psi(0), \delta_1) \subset \Psi(\Delta(0, r))$. For each $w \in \Delta(\Psi(0), \delta_1)$, we have $|\Psi(z) - w| > \delta_1$ on $\Gamma(0, r)$, and we have for sufficiently large n ,

$$|(\Psi_n(z) - w) - (\Psi(z) - w)| = |\Psi_n(z) - \Psi(z)| < \delta_1, \quad z \in \Gamma(0, r).$$

By the argument principle, for sufficiently large n ,

$$n \left(r, \frac{1}{\Psi_n(z) - w} \right) = n \left(r, \frac{1}{\Psi(z) - w} \right) = 1.$$

This shows that for each $w \in \Delta(\Psi(0), \delta_1)$, there exists $z_0 \in \Delta(0, r)$ such that $\Psi_n(z_0) = w$. Hence (b) holds.

Obviously, there exists δ_2 such that $\Psi(\Delta(0, \delta_2)) \subset \Delta(\Psi(0), \delta_1/2)$, and hence $\Psi(\Delta(0, \delta_2)) \subset \Delta(\Psi(0), \delta_1)$. Since $\Psi_n(z) \Rightarrow \Psi(z)$ in $\Delta(0, R)$, we have that for sufficiently large n , $|\Psi_n(z) - \Psi(z)| < \delta_1/2$ in $\Delta(0, \delta_2)$ and hence

$$|\Psi_n(z) - \Psi(0)| \leq |\Psi_n(z) - \Psi(z)| + |\Psi(z) - \Psi(0)| < \delta_1/2 + \delta_1/2 = \delta_1$$

in $\Delta(0, \delta_2)$. Hence (c) holds. $\square$

Lemma 3.8. *Let $\{f_n\}$ be a sequence of meromorphic functions in $\Delta(z_0, r)$, all of whose zeros are multiple, and let $\{\psi_n\}$ be a sequence of meromorphic functions in $\Delta(z_0, r)$ such that $\psi_n \Rightarrow \psi$ in $\Delta(z_0, r)$, where ψ is a non-vanishing holomorphic function in*

$\Delta(z_0, r)$. Let E be a (countable) discrete set in $\Delta(z_0, r)$ which has no accumulation points in $\Delta(z_0, r)$. Suppose that

- (a) $f_n(z) \xrightarrow{X} f(z)$ in $\Delta(z_0, r) \setminus E$;
- (b) for some $a_1 \in E$, no subsequence of $\{f_n\}$ is normal at a_1 ;
- (c) for all $n \in \mathbb{N}$, $f'_n(z) \neq \psi_n(z)$ in $\Delta(z_0, r)$.

Then

- (d) There exists $r \geq 0$ such that for sufficiently large n , f_n has a single zero $z_{n,1}$ of order 2 and a single pole $z_{n,2}$ of order 1 in $\Delta(a_1, r)$, where $z_{n,i} \rightarrow a_1$ as $n \rightarrow \infty$, $i = 1, 2$;
- (e) $f(z) = \int_{a_1}^z \psi(\zeta) d\zeta$.

Proof. Without loss of generality, we may suppose that $a_1 = 0$. Set $\Psi(z) = \int_{\zeta=0}^z \psi(\zeta) d\zeta$. There exists $\frac{1}{2} > \delta > 0$ such that $\Psi(z)$ is a univalent function in $\Delta(0, 2\delta)$. Since $\psi_n(z) \Rightarrow \psi(z)$ on Δ , without loss of generality, we may suppose that for all $n \in \mathbb{N}$, ψ_n is a holomorphic function in $\Delta(0, 2\delta)$.

Let $\Psi_n(z) = \int_{\zeta=0}^z \psi_n(\zeta) d\zeta$, $z \in \Delta(0, 2\delta)$. Obviously, we have that $\Psi_n(z) \Rightarrow \Psi(z)$ in $\Delta(0, 2\delta)$. By Lemma 3.7, we have for sufficiently large n ,

- (a) $\Psi_n(z)$ is a holomorphic univalent function in $\Delta(0, \delta)$;
- (b) there exists $\delta_1 > 0$ such that $\Delta(\Psi(0), \delta_1) \subset \Psi(\Delta(0, \delta))$ and $\Delta(\Psi(0), \delta_1) \subset \Psi_n(\Delta(0, \delta))$;
- (c) there exists $\delta_2 > 0$ such that $\Psi(\Delta(0, \delta_2)) \subset \Delta(\Psi(0), \delta_1)$ and $\Psi_n(\Delta(0, \delta_2)) \subset \Delta(\Psi(0), \delta_1)$.

For convenience, we suppose that for all n , (a), (b) and (c) hold.

Now we consider $\Psi(z)$ and $\Psi_n(z)$ only in $\Delta(0, \delta)$. Let $F_n(w) = f_n(\Psi_n^{-1}(w))$, $w \in \Delta(\Psi(0), \delta_1)$. Then $f_n(z) = F_n(\Psi_n(z))$ in $\Delta(0, \delta_2)$.

We claim that no subsequence of $\{F_n(w)\}$ is normal at $\Psi(0)$. Otherwise, suppose that $\{F_n(w)\}$ is normal at $\Psi(0)$ (without loss of generality, we call also the subsequence $\{F_n(w)\}$). Since $\Psi_n(z) \Rightarrow \Psi(z)$ in $\Delta(0, 2\delta)$, $\{F_n(\Psi_n(z))\}$ is normal at 0. i.e., $\{f_n(z)\}$ is normal at 0, a contradiction.

Now we have

$$F'_n(w) = f'_n(\Psi_n^{-1}(w)) \frac{1}{\Psi'_n(z)} = f'_n(z) \frac{1}{\psi_n(z)} \neq \psi_n(z) \frac{1}{\psi_n(z)} = 1$$

in $\Delta(\Psi(0), \delta_1)$. Obviously, all the zeros of $F_n(w)$ in $\Delta(\Psi(0), \delta_1)$ are multiple. Clearly, $\{F_n\}_{n=1}^\infty$ is normal in $\Delta'(\psi(0), \delta_1)$. Then by Lemma 2.9, there exists $0 < \delta_3 < \delta_1$ such that $F_n(w)$ has only a single (multiple) zero in $\Delta(\Psi(0), \delta_3)$ for sufficiently large n . Do as in Lemma 3.7, we have that for sufficiently large n , there exists $\delta_4 > 0$ such that $\Psi_n(\Delta(0, \delta_4)) \subset \Delta(\Psi(0), \delta_3)$. Therefore, $f_n(z) = F_n(\Psi_n(z))$ has at most a single (multiple) zero in $\Delta(0, \delta_4)$.

We claim that for sufficiently large n , f_n has a single zero $z_{n,1}$ of order 2 in $\Delta(0, \delta_4)$, where $z_{n,1} \rightarrow 0$ as $n \rightarrow \infty$. It suffices to prove that each subsequence of $\{f_n\}$ has a subsequence $\{f_m\}$ such that, for sufficiently large m , f_m has a single zero $z_{m,1}$ of order

2 in $\Delta(0, \delta_4)$, where $z_{m,1} \rightarrow 0$ as $m \rightarrow \infty$. Suppose that we have a subsequence of $\{f_n\}$, which (to avoid complication in notion) we again call $\{f_n\}$. Since $\{f_n\}$ is not normal at 0, it follows from Lemma 2.5 that we can extract a subsequence $\{f_m\}$ of $\{f_n\}$, points $z_m \rightarrow 0$, and positive numbers $\rho_m \rightarrow 0$ such that

$$g_m(\zeta) = \frac{f_m(z_m + \rho_m \zeta)}{\rho_m} \xrightarrow{\chi} g(\zeta), \quad \zeta \in \mathbb{C},$$

where g is a nonconstant meromorphic function of finite order on $\mathbb{C}$, all of whose zeros are multiple. Now we have

$$0 \neq f'_m(z_m + \rho_m \zeta) - \psi_m(z_m + \rho_m \zeta) \Rightarrow g'(\zeta) - \psi(0), \quad \zeta \in \mathbb{C} - \{g^{-1}(\infty)\}.$$

By Hurwitz's Theorem, either $g'(\zeta) \neq \psi(0)$ on $\mathbb{C}$, or $g'(\zeta) \equiv \psi(0)$ on $\mathbb{C}$. In the latter case, $g(\zeta) = \psi(0)\zeta + c$, which contradicts the fact that all zeros of g are multiple. Thus $g'(\zeta) \neq \psi(0)$ on $\mathbb{C}$. By Lemma 2.4, $g(\zeta) = \frac{\psi(0)(\zeta-a)^2}{\zeta-b}$ for distinct complex numbers a and b . It now follows from Hurwitz's Theorem that there exist sequence $\zeta_{m,1} \rightarrow a$ such that, for sufficiently large m , $g_m(\zeta_{m,1}) = 0$. Obviously, $\zeta_{m,1}$ is a zero of order 2 of $g_m(\zeta)$. Set $z_{m,1} = z_m + \rho_m \zeta_{m,1}$, we have $z_{m,1} \rightarrow 0$ and $f_m(z_{m,1}) = 0$. Obviously $z_{m,1}$ is a zero of order 2 of $f_m(z)$ and since we have proved that there is at most one such zero in $\Delta(0, \delta_4)$ (for sufficiently large m), it is the only one, as required.

By Lemma 2.8, there exists $\eta > 0$ such that f_n has a single simple pole $z_{n,2}$ in $\Delta(0, \eta)$ for sufficiently large n . We claim that $z_{n,2} \rightarrow 0$ as $n \rightarrow \infty$. Otherwise, there exist $0 < \delta_5 < \delta_4$ and a subsequence of $\{f_n\}$ (that we continue to call $\{f_n\}$) such that $f_n(z) \neq \infty$, $z \in \Delta(0, \delta_5)$. By Lemma 2.6, $\{f_n\}$ is normal at 0. A contradiction. Set $r = \min\{\delta_4, \eta\}$. Hence (d) holds.

By Lemma 2.8, we have $f(z) = \int_{a_1}^z \psi(\zeta) d\zeta$. Hence (e) holds. $\square$

Lemma 3.9. (cf. [14, Lemma 7]) *Let $\{f_n\}$ be a family of meromorphic functions on the plane domain D , all of whose zeros are multiple, and let $\{\psi_n\}$ be a sequence of meromorphic functions in D such that $\psi_n \xrightarrow{\chi} \psi$ in D , where $\psi(z) \not\equiv 0, \infty$ in D . If for each $n \in \mathbb{N}$, $f'_n \neq \psi_n$ in D , then $\{f_n\}$ is quasinormal in D .*

Proof. It suffices to show that $\{f_n\}$ is quasinormal in a neighborhood of each point of D . Let $p \in D$. There exists $t > 0$ such that $\Delta(p, t) \subset D$ and ψ is holomorphic and does not vanish in $\Delta'(p, t)$.

For each $q \in \Delta'(p, t)$, Let $\Psi(z) = \int_{\zeta=q}^z \psi(\zeta) d\zeta$ in $\Delta'(p, t)$. Since $\psi(z)$ is holomorphic and does not vanish in $\Delta'(p, t)$ and $\psi_n \xrightarrow{\chi} \psi$ in D , there exists $0 < R < t - |p - q|$ such that for sufficiently large n , $\Psi(z)$ is a holomorphic univalent function in $\Delta(q, R)$ and $\psi_n(z)$ is a holomorphic function in $\Delta(q, R)$. Let $\Psi_n(z) = \int_{\zeta=q}^z \psi_n(\zeta) d\zeta$ in $\Delta(q, R)$. Obviously, we have $\Psi_n(z) \Rightarrow \Psi(z)$ in $\Delta(q, R)$.

Let $0 < r < R$. By Lemma 3.7, we have, for sufficiently large n ,

- (a) $\Psi_n(z)$ is a holomorphic univalent function in $\Delta(q, r)$;
- (b) there exists δ_1 such that $\Delta(\Psi(q), \delta_1) \subset \Psi(\Delta(q, r))$ and $\Delta(\Psi(q), \delta_1) \subset \Psi_n(\Delta(q, r))$;
- (c) there exists δ_2 such that $\Psi(\Delta(q, \delta_2)) \subset \Delta(\Psi(q), \delta_1)$ and $\Psi_n(\Delta(q, \delta_2)) \subset \Delta(\Psi(q), \delta_1)$.

For convenience in notation, we assume that for all n , (a), (b) and (c) hold.

Now we consider $\Psi(z)$ and $\Psi_n(z)$ only in $\Delta(q, r)$. Let $F_n(w) = f_n(\Psi_n^{-1}(w))$, $w \in \Delta(\Psi(q), \delta_1)$. Then $f_n(z) = F_n(\Psi_n(z))$ in $\Delta(q, \delta_2)$.

Now we have

$$F'_n(w) = f'_n(\Psi_n^{-1}(w)) \frac{1}{\Psi'_n(z)} = f'_n(z) \frac{1}{\psi_n(z)} \neq \psi_n(z) \frac{1}{\psi_n(z)} = 1$$

in $\Delta(\Psi(q), \delta_1)$. Obviously all the zeros of $F_n(w)$ are multiple in $\Delta(\Psi(q), \delta_1)$. By Lemma 2.10, $\{F_n(w)\}$ is quasinormal in $\Delta(\Psi(q), \delta_1)$. Since $f_n(z) = F_n(\psi_n(z))$ in $\Delta(q, \delta_2)$ and $\psi_n(z) \Rightarrow \psi(z)$ in $\Delta(q, R)$, $\{f_n(z)\}$ is quasinormal in $\Delta(q, \delta_2)$ and hence quasinormal in $\Delta'(p, t)$.

Suppose now that $\{f_n\}$ is not quasinormal at p . Then there exists points $z_j \in \Delta'(p, \delta)$ ($j = 1, 2, \dots$) and a subsequence of $\{f_n\}$ (that we continue to call $\{f_n\}$) such that $z_j \rightarrow p$ as $j \rightarrow \infty$ and no subsequence of $\{f_n\}$ is normal at any z_j , $j = 1, 2, \dots$, see [11, Thm. 4.4]. Let $E = \{z_j : j = 1, 2, \dots\}$. Taking a subsequence of $\{f_n\}$ (that we continue to call $\{f_n\}$), we may assume that $f_n \xrightarrow{\lambda} H$ in $\Delta'(p, \delta) \setminus E$. By Lemma 3.8, we have $H' \equiv \psi$ and $H(z_j) = 0$. It follows that H is holomorphic in $\Delta'(p, t)$ and $H' \equiv \psi$ there. Moreover, since ψ has no essential singularity at p , the same is true of H . But then $H(z_j) = 0$ for $j = 1, 2, \dots$ implies $H \equiv 0$, which contradict $H' \equiv \psi \neq 0$. $\square$

Lemma 3.10. *Let $\{f_n\}$ be a family of meromorphic functions in $\Delta(z_0, r)$ and $k \neq 0$ an integer. Let $\{b_n\}$ be a sequence of holomorphic functions in $\Delta(z_0, r)$ such that $b_n \Rightarrow b$ in $\Delta(z_0, r)$, where $b(z) \neq 0$ is a holomorphic function in $\Delta(z_0, r)$. Suppose that*

$$f_n(z) \neq 0, f'_n(z) \neq (z - z_0)^k b_n(z), \quad z \in \Delta(z_0, r).$$

Then $\{f_n\}$ is normal in $\Delta(z_0, r)$.

Proof. Without loss of generality, we assume that $z_0 = 0$ and $r = 1$. By Lemma 2.7, $\{f_n\}_{n=1}^\infty$ is normal in Δ' . Without loss of generality, $f_n \xrightarrow{\lambda} f$ in Δ' , and thus since $f_n \neq 0$, then $\frac{1}{f_n} \Rightarrow \frac{1}{f}$ in Δ' . The possibilities of the limit function $\frac{1}{f}$ are $\frac{1}{f} \equiv \infty$ or that $\frac{1}{f}$ is a holomorphic function. In the last case, we get by the maximum principle that $\{\frac{1}{f_n}\}_{n=1}^\infty$ converges to some holomorphic function in Δ and we are done. Hence we can assume that $\frac{1}{f} \equiv \infty$, i.e., $f \equiv 0$ in Δ' .

Obviously, we have

$$f'_n(z) - z^k b_n(z) \Rightarrow -z^k b(z), \quad \text{in } \Delta'.$$

Since $f'_n(z) - z^k b_n(z) \neq 0$ in Δ , we deduce that

$$H_n(z) = \frac{1}{f'_n(z) - z^k b_n(z)}$$

is holomorphic in Δ , so by the maximum principle

$$(3.4) \quad H_n(z) \Rightarrow -\frac{1}{z^k b(z)}, \quad z \in \Delta.$$

Obviously, $-\frac{1}{z^k b(z)}$ is an analytic function in Δ and hence k is a negative integer. Let $m = -k$. By (3.4) and Rouché's Theorem for holomorphic function, for sufficiently

large n ,

$$(3.5) \quad n \left(\frac{1}{2}, f'_n(z) - \frac{b_n(z)}{z^m} \right) = n \left(\frac{1}{2}, \frac{1}{H_n(z)} \right) = n \left(\frac{1}{2}, -\frac{b(z)}{z^m} \right) = m.$$

Since $f'_n(z) \neq \frac{b_n(z)}{z^m}$, then $f'_n(0) \neq \infty$ and hence $f_n(0) \neq \infty$. Obviously, 0 is a pole of $f'_n(z) - \frac{b_n(z)}{z^m}$ of order m . By (3.5), for sufficiently large n , f_n has no poles in $\Delta(0, \frac{1}{2})$, i.e., f_n is holomorphic in $\Delta(0, \frac{1}{2})$. We have proved that $f_n \Rightarrow 0$ in $\Delta'(z_0, r)$, so, by the maximum principle, $f_n \Rightarrow 0$ in $\Delta(0, \frac{1}{2})$ and $\{f_n\}$ is normal at 0. $\square$

Lemma 3.11. *Let $\mathcal{F} = \{f_n\}$ be a family of holomorphic functions in $\Delta(z_0, r)$, all of whose zeros are multiple. Let $k \geq 2$ be a positive integer and $\{b_n\}$ be a sequence of holomorphic functions in $\Delta(z_0, r)$ such that $b_n \Rightarrow b$ in $\Delta(z_0, r)$, where $b(z) \neq 0$ is a holomorphic function in $\Delta(z_0, r)$. Suppose that*

$$f'_n(z) \neq \frac{b_n(z)}{z^k}, \quad z \in \Delta(z_0, r).$$

Then $\mathcal{F}$ is normal in $\Delta(z_0, r)$.

Proof. Without loss of generality, we assume that $z_0 = 0$ and $r = 1$. By Lemma 2.6, $\mathcal{F}$ is normal in Δ' . Suppose that $\mathcal{F}$ is not normal at 0. Let $\mathcal{F}_1 = \{F_n\}$, where $F_n(z) = z^k f_n(z)$. Obviously, all zeros of $\mathcal{F}_1$ are multiple in Δ . By Lemma 3.3, $\mathcal{F}_1$ is not normal at 0. By Lemma 2.5, there exist points $z_n \rightarrow 0$, and positive numbers $\rho_n \rightarrow 0$ and a subsequence of $\{F_n\}$ (that we continue to call $\{F_n\}$) such that

$$(3.6) \quad g_n(\zeta) = \frac{F_n(z_n + \rho_n \zeta)}{\rho_n} = \frac{(z_n + \rho_n \zeta)^k f_n(z_n + \rho_n \zeta)}{\rho_n} \Rightarrow g(\zeta), \quad \zeta \in \mathbb{C},$$

where $g(\zeta)$ is a nonconstant holomorphic function in $\mathbb{C}$, all of whose zeros are multiple.

We consider the following two cases.

Case 1: $z_n/\rho_n \rightarrow \infty$.

Observe first that since $\frac{z}{z_n + \rho_n \zeta} \Rightarrow 1$ in $\mathbb{C}$, then (3.6) is equivalent to $\frac{z_n^k}{\rho_n} f_n(z_n + \rho_n \zeta) \Rightarrow g(\zeta)$ in $\mathbb{C}$. Differentiating gives $z_n^k f'_n(z_n + \rho_n \zeta) \Rightarrow g'(\zeta)$ in $\mathbb{C}$, and by the same reasoning as above it is equivalent to $(z_n + \rho_n \zeta)^k f'_n(z_n + \rho_n \zeta) \Rightarrow g'(\zeta)$ in $\mathbb{C}$.

Now we have

$$0 \neq \frac{f'_n(z_n + \rho_n \zeta)}{\frac{b_n(z_n + \rho_n \zeta)}{(z_n + \rho_n \zeta)^k}} - 1 = \frac{(z_n + \rho_n \zeta)^k f'_n(z_n + \rho_n \zeta)}{b_n(z_n + \rho_n \zeta)} - 1 \Rightarrow \frac{g'(\zeta)}{b(0)} - 1 \quad \text{in } \mathbb{C}.$$

By Hurwitz's Theorem, either $g'(\zeta) - b(0) \equiv 0$ in $\mathbb{C}$, or $g'(\zeta) - b(0) \neq 0$ in $\mathbb{C}$. If $g'(\zeta) - b(0) \equiv 0$, then $g(\zeta) = b(0)\zeta + c$ which is impossible since all the zeros of $g(\zeta)$ must be multiple. If $g'(\zeta) - b(0) \neq 0$, then, by Lemma 2.4, $g(\zeta) = \frac{b(0)(\zeta - a)^2}{\zeta - b}$ which contradicts that $g(\zeta)$ is an entire function.

Case 2: $z_n/\rho_n \rightarrow \alpha$ in $\mathbb{C}$

We have

$$G_n(\zeta) = \frac{F_n(\rho_n \zeta)}{\rho_n} = \frac{F_n(z_n + \rho_n(\zeta - \frac{z_n}{\rho_n}))}{\rho_n} \Rightarrow g(\zeta - \alpha) \quad \text{in } \mathbb{C}.$$

Writing

$$G(\zeta) = g(\zeta - \alpha) \quad \text{and} \quad \psi_n(\zeta) = \rho_n^{k-1} f_n(\rho_n \zeta) = \frac{F_n(\rho_n \zeta)}{\rho_n} \frac{1}{\zeta^k},$$

we have

$$\psi_n(\zeta) \Rightarrow \frac{G(\zeta)}{\zeta^k} = \psi(\zeta), \quad \text{in } \mathbb{C} \setminus \{0\}.$$

Since for all n , $\psi_n(\zeta)$ is a holomorphic function, we have, by the maximum principle,

$$\psi_n(\zeta) \Rightarrow \psi(\zeta) \quad \text{in } \mathbb{C},$$

where $\psi(\zeta)$ is an entire function. Hence we have

$$0 \neq \rho_n^k \left(f'_n(\rho_n \zeta) - \frac{b_n(\rho_n \zeta)}{(\rho_n \zeta)^k} \right) = \psi'_n(\zeta) - \frac{b_n(\rho_n \zeta)}{\zeta^k} \Rightarrow \psi'(\zeta) - \frac{b(0)}{\zeta^k} \quad \text{in } \mathbb{C} \setminus \{0\}.$$

By Hurwitz's Theorem, either $\psi'(\zeta) \equiv \frac{b(0)}{\zeta^k}$ in $\mathbb{C} \setminus \{0\}$ (and hence in $\mathbb{C}$), or $\psi'(\zeta) \neq \frac{b(0)}{\zeta^k}$ in $\mathbb{C} \setminus \{0\}$ (and in fact in $\mathbb{C}$, since $b(0)/\zeta^k = \infty$ at $\zeta = 0$). Since ψ is a holomorphic function, the first alternative obviously cannot hold. Thus $\psi'(\zeta) \neq \frac{b(0)}{\zeta^k}$. It then follows from Theorem E and Lemma 3.1 that $\psi(\zeta) \equiv c$. Since g is not a constant function and $g(\zeta - \alpha) = \zeta^k \psi(\zeta)$, we have $c \neq 0$. Now we have

$$(3.7) \quad \psi_n(\zeta) = \rho_n^{k-1} f_n(\rho_n \zeta) \Rightarrow c, \quad \text{in } \mathbb{C}.$$

Now suppose that there exists $\delta > 0$ such that for sufficiently large n , $f_n(z) \neq 0$ for $z \in \Delta(0, \delta)$. Then by Lemma 3.10, $\{f_n\}$ is normal at 0. A contradiction.

Otherwise, taking a subsequence and renumbering if necessary, we may assume that there exist $z_n^* \rightarrow 0$ such that $f_n(z_n^*) = 0$. We may assume that z_n^* is the zero of f_n of smallest modulus. By (3.7), we have, for sufficiently large n , $f_n(0) \neq 0$ and hence $z_n^* \neq 0$. By Hurwitz's Theorem and (3.7), we have $\frac{z_n^*}{\rho_n} \rightarrow \infty$.

Let $G_n(\zeta) = (z_n^*)^{k-1} f_n(z_n^* \zeta)$. Then we have

$$G'_n(\zeta) = (z_n^*)^k f'_n(z_n^* \zeta) \neq \frac{b(z_n^* \zeta)}{\zeta^k}.$$

Since $G_n(\zeta) \neq 0$ in Δ , it follows from Lemma 3.10 that $\{G_n\}$ is normal in Δ . By Lemma 3.9, $\{G_n\}$ is quasinormal in $\mathbb{C}$. Thus, there exists a subsequence of $\{G_n\}$ (that we continue to call $\{G_n\}$) and $E \subset \mathbb{C}$ such that

- (b1) E have no accumulation point in $\mathbb{C}$;
- (b2) $G_n(\zeta) \xrightarrow{x} G(\zeta)$ in $\mathbb{C} \setminus E$;
- (b3) for each $\zeta_0 \in E$, no subsequence of $\{G_n\}$ is normal at ζ_0 .

Obviously, we have $E \cap \Delta = \emptyset$ and all zeros of $G(\zeta)$ are multiple.

Since $G_n(0) = \left(\frac{z_n^*}{\rho_n}\right)^{k-1} \psi_n(0) \rightarrow \infty$ and $\{G_n(\zeta)\}$ is a family of holomorphic functions, we have $G(\zeta) \equiv \infty$ in $\mathbb{C} \setminus E$. Suppose that $1 \notin E$. Since $G_n(1) = 0$, we have $G(1) = 0$ which contradicts that $G(\zeta) \equiv \infty$ on $\mathbb{C} \setminus E$. Thus we have $1 \in E$. By Lemma 3.8, $G(\zeta) = \int_1^\zeta \frac{b(0)}{\xi^k} d\xi$ on $\mathbb{C} \setminus E$, which also contradicts that $G(\zeta) \equiv \infty$ in $\mathbb{C} \setminus E$. $\square$

4. PROOF OF THEOREM

Proof. We assume that $f' = h$ has at most finitely many zeros and derive a contradiction.

We claim that there exist $t_n \rightarrow \infty$ and $\varepsilon_n \rightarrow 0$ such that

$$(4.1) \quad S(\Delta(t_n, \varepsilon_n), f) = \frac{1}{\pi} \iint_{|z-t_n| < \varepsilon_n} [f^\#(z)]^2 dx dy \rightarrow \infty.$$

Otherwise there would exist $\varepsilon > 0$ and $M > 0$ such that for all $z_0 \in \mathbb{C}$, we have

$$S(\Delta(z_0, \varepsilon), f) = \frac{1}{\pi} \iint_{|z-z_0| < \varepsilon} [f^\#(z)]^2 dx dy < M.$$

Then $S(r, f) = \frac{1}{\pi} \iint_{|z| < r} [f^\#(z)]^2 dx dy = O(r^2)$. Thus

$$T_0(r, f) = \int_0^r \frac{S(t)}{t} dt = O(r^2).$$

On the other hand, by Lemma 2.2, $T_0(r, h) = Ar^2(1 + (1))$ as $r \rightarrow \infty$ where $A > 0$ is a constant. Hence it is impossible that $T_0(r, h) = o\{T_0(r, f)\}$ and (4.1) follows.

Let ω_1, ω_2 be the two fundamental periods of $h(z)$ and $P(0 \in P)$ be a fundamental parallelogram of $h(z)$. There exist integers i_n and j_n such that $z_n \in P$, where $z_n = t_n - i_n\omega_1 - j_n\omega_2$. There exists a subsequence of $\{z_n\}$ (that we continue to call $\{z_n\}$) such that $z_n \rightarrow z_0$ as $n \rightarrow \infty$. Let $f_n(z) = f(z + i_n\omega_1 + j_n\omega_2)$. By (4.1) we have

$$(4.2) \quad S(\Delta(z_n, \varepsilon_n), f_n) = S(\Delta(t_n, \varepsilon_n), f) \rightarrow \infty,$$

and hence, there exists z_n^* ($z_n^* \rightarrow z_0$) such that $f_n^\#(z_n^*) \rightarrow \infty$ as $n \rightarrow \infty$. Without loss of generality, we can assume that $z_0 = 0$. Hence we have that no subsequence of $\{f_n\}$ is normal at 0 and by (4.2),

$$(4.3) \quad S(\Delta(z_n, \varepsilon_n), f_n) \rightarrow \infty,$$

where $z_n \rightarrow 0$ and $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$.

There exists $R > 0$ such that $\overline{P} \subset \Delta(0, R)$ and $\Delta(z_n, \varepsilon_n) \subset \Delta(0, R)$ for each n . Set $D = \Delta(0, R)$. Obviously, we have $z_0 \in D$. By assumption, for sufficiently large n ,

$$f'_n(z) = f'(z + i_n\omega_1 + j_n\omega_2) \neq h(z + i_n\omega_1 + j_n\omega_2) = h(z), \quad z \in D.$$

Without loss of generality, we can assume that for all $n \in \mathbb{N}$, $f'_n(z) \neq h(z)$ in D .

Now, $\{f_n\}$ is a family of functions meromorphic in D such that

- (a*) all zeros of $\{f_n\}$ are multiple in D ;
- (b*) for each n , $f'_n(z) \neq h(z)$ in D , where h is a nonconstant elliptic function;
- (c*) no subsequence of $\{f_n\}$ is normal at 0.

It follows from Lemma 3.9 that $\{f_n\}$ is quasinormal in D . Hence there exists $\tau > 0$ such that $\{f_n\}$ is normal in $\Delta'(0, \tau)$ and $h(z) \neq 0$ on $\Delta'(0, \tau)$.

Without loss of generality, we may assume that $\tau = 1$. Then there exists a subsequence of $\{f_n\}$ (that we continue to call $\{f_n\}$) such that

- (a) all zeros of $\{f_n\}$ are multiple in Δ ;
- (b) for each n , $f'_n(z) \neq h(z)$ in Δ , where $h \neq 0$ in Δ' ;

(c) no subsequence of $\{f_n\}$ is normal at 0.

(d) $f_n(z) \Rightarrow f_0(z)$ on $\Delta'(0, 1)$

Case 1: $h(0) \neq 0, \infty$.

By Lemma 3.8, $f_0(z) = \int_{\zeta=0}^z h(\zeta)d\zeta$ and for sufficiently large n , there exists $1 > \delta > 0$ such that f_n has a single zero of order 2 in $\Delta(0, \delta)$. By Lemma 3.6, there exists $M > 0$ such that $S(\frac{\delta}{2}, f_n) < M$ which contradicts (4.3).

Case 2: $h(0) = 0$.

Suppose that 0 is a zero of order k of $h(z)$, where k is a positive integer. Let us assume, making standard normalizations, that for $z \in \Delta$

$$h(z) = z^k + a_{k+1}z^{k+1} + \dots = z^k \hat{h}(z)$$

where $\hat{h}(z) \neq 0, \infty$ in $\Delta(0, 1)$ and $\hat{h}(0) = 1$.

We claim that for each $\delta > 0$, there exists at least one zero of f_n in $\Delta(0, \delta)$ for sufficiently large n . Otherwise, there exists a subsequence of $\{f_n\}$ (that we continue to call $\{f_n\}$) such that $f_n(z) \neq 0$ in $\Delta(0, \delta)$. By Lemma 2.3, $\{f_n\}$ is normal at 0. A contradiction.

Hence, taking a subsequence and renumbering if necessary, we may assume that $a_n \rightarrow 0$ is the zero of f_n of smallest modulus. Since $f'_n(z) \neq h(z)$ and all the zeros of $\{f_n\}$ are multiple, we have $f_n(0) \neq 0$ and hence $a_n \neq 0$. Let $F_n = \frac{f_n(a_n \zeta)}{a_n^{k+1}}$. We have that

- (a1) $F_n(\zeta) \neq 0$ in Δ ;
- (a2) all zeros of $F_n(\zeta)$ are multiple;
- (a3) $F'_n(\zeta) \neq \zeta^k \hat{h}(a_n \zeta)$ and $F_n(1) = 0$.

By Lemma 3.10, $\{F_n(\zeta)\}$ is normal in Δ . By Lemma 3.9, $\{F_n(\zeta)\}$ is quasinormal on $\mathbb{C}$. Thus, there exists a subsequence of $\{F_n(\zeta)\}$ (that we continue to call $\{F_n(\zeta)\}$) and $E_1 \subset \mathbb{C}$ such that

- (b1) E_1 have no accumulation point in $\mathbb{C}$;
- (b2) $F_n(\zeta) \xrightarrow{x} F(\zeta)$ on $\mathbb{C} \setminus E_1$;
- (b3) for each $\zeta_0 \in E_1$, no subsequence of $\{F_n(\zeta)\}$ is normal at ζ_0 .

Obviously, $E_1 \cap \Delta = \emptyset$ and all zeros of $F(\zeta)$ are multiple in $\mathbb{C} \setminus E_1$.

Case 2.1: $1 \notin E_1$

Since all zeros of $\{F_n(\zeta)\}$ are multiple and $F_n(1) = 0$, we have that $F(1) = F'(1) = 0$ (recall that $\hat{h}(0) = 1$) and hence $F(\zeta)$ is a meromorphic function in $\mathbb{C} \setminus E_1$.

We claim that $E_1 = \emptyset$. Otherwise, let $\zeta_0 \in E_1$. Obviously, $\zeta_0 \neq 0$. By Lemma 3.8, $F'(\zeta) = \zeta^k$. Recall that $\hat{h}(0) = 1$ and hence $F'(1) = 1$ which contradicts that $F'(1) = 0$.

By Lemma 3.4, either $F'(\zeta) \equiv \zeta^k$ in $\mathbb{C}$, or $F'(\zeta) \neq \zeta^k$ in $\mathbb{C}$. If $F'(\zeta) \equiv \zeta^k$ in $\mathbb{C}$, then $F'(1) = 1$ which contradicts that $F'(1) = 0$. If $F'(\zeta) \neq \zeta^k$ on $\mathbb{C}$. By Theorem E, F

must be rational and then by Lemma 3.2,

$$F(\zeta) = \frac{\prod_{i=1}^{m+k+1} (\zeta - \alpha_i)}{(k+1)(\zeta - \beta)^m},$$

where m is a nonnegative integer, $\beta \in \mathbb{C}$, $\alpha_i \neq 0, \beta$, $1 \leq i \leq m+k+1$.

Hence, we have

$$(4.4) \quad F_n(\zeta) \xrightarrow{\chi} \frac{\prod_{i=1}^{m+k+1} (\zeta - \alpha_i)}{(k+1)(\zeta - \beta)^m}, \quad \text{in } \mathbb{C}$$

By Hurwitz's Theorem, there exist sequences $\zeta_{n,i} \rightarrow \alpha_i$ and $\eta_{n,j} \rightarrow \beta$ as $n \rightarrow \infty$ (counting multiplicities of zeros and poles, respectively), such that for sufficiently large n , $F_n(\zeta_{n,i}) = 0$ and $F_n(\eta_{n,j}) = \infty$, where $i = 1, 2, \dots, m+k+1$ and $j = 1, 2, \dots, m$. Writing $z_{n,i} = a_n \zeta_{n,i}$. Thus, $f_n(z_{n,i}) = 0$ and $z_{n,i} \rightarrow 0$ as $n \rightarrow \infty$, where $i = 1, 2, \dots, m+k+1$. Set $B_n = \{z_{n,1}, z_{n,2}, \dots, z_{n,m+k+1}\}$.

We claim that for each $\delta > 0$, there exists at least $m+k+2$ zeros of f_n in $\Delta(0, \delta)$ for sufficiently large n . Otherwise, there exists a subsequence of $\{f_n\}$ (that we continue to call $\{f_n\}$) such that $f_n(z)$ have $m+k+1$ zeros in $\Delta(0, \delta)$. If $f_0(z) \equiv \infty$, then by Lemma 3.6, there exists $M > 0$ such that for n sufficiently large, $S(\frac{\delta}{2}, f_n) < M$ which contradicts (4.3). If $f_0(z) \not\equiv \infty$, then by Lemma 3.5, $f_0(z) = \frac{\zeta^{k+1+c}}{k+1}$ in Δ' , where c is a constant. By Lemma 3.6, there exists $M > 0$ such that for n sufficiently large, $S(\frac{\delta}{2}, f_n) < M$ which also contradicts (4.3).

Hence, taking a subsequence and renumbering if necessary, we may assume that $b_n \rightarrow 0$ is the zero of f_n of smallest modulus in $\Delta \setminus B_n$. Set $r_n = \frac{a_n}{b_n}$. Then we have $F_n(\frac{1}{r_n}) = 0$. Since $b_n \notin B_n$, Obviously $\frac{1}{r_n} \neq \zeta_{n,i}$, where $i = 1, 2, \dots, m+k+1$. By Hurwitz's Theorem and (4.4), we have $\frac{1}{r_n} \rightarrow \infty$ and hence $r_n \rightarrow 0$ as $n \rightarrow \infty$.

Let $G_n(\zeta) = \frac{f_n(b_n \zeta)}{b_n^{k+1}}$. We have that for sufficiently large n ,

- (d1) $G_n(\zeta)$ have only $m+k+1$ zeros $r_n \zeta_{n,i}$ in Δ . Obviously, $|r_n \zeta_{n,i}| \rightarrow 0$, as $n \rightarrow \infty$, where $i = 1, 2, \dots, m+k+1$;
- (d2) all zeros of $G_n(\zeta)$ are multiple;
- (d3) $G'_n(\zeta) \neq \zeta^k \widehat{h}(b_n \zeta)$ and $G_n(1) = 0$.

By Lemma 3.10, $\{G_n(\zeta)\}$ is normal in Δ' . By Lemma 3.9, $\{G_n(\zeta)\}$ is quasinormal on $\mathbb{C}$. Thus, there exists a subsequence of $\{G_n(\zeta)\}$ (that we continue to call $\{G_n(\zeta)\}$) and $E_2 \subset \mathbb{C}$ such that

- (e1) E_2 have no accumulation point in $\mathbb{C}$;
- (e2) $G_n(\zeta) \xrightarrow{\chi} G(\zeta)$ on $\mathbb{C} \setminus E_2$;
- (e3) for each $\zeta_0 \in E_2$, no subsequence of $\{G_n(\zeta)\}$ is normal at ζ_0 .

Obviously, $E_2 \cap \Delta' = \emptyset$ and all zeros of $G(\zeta)$ are multiple in $\mathbb{C} \setminus E_2$.

Let

$$G_n^*(\zeta) = G_n(\zeta) \frac{\prod_{j=1}^m (\zeta - r_n \eta_{n,j})}{\prod_{i=1}^{m+k+1} (\zeta - r_n \zeta_{n,i})},$$

$$F_n^*(\zeta) = F_n(\zeta) \frac{\prod_{j=1}^m (\zeta - \eta_{n,j})}{\prod_{i=1}^{m+k+1} (\zeta - \zeta_{n,i})}.$$

By (4.4),

$$G_n^*(r_n \zeta) = F_n^*(\zeta) \Rightarrow \frac{1}{k+1}, \quad \zeta \in \mathbb{C}$$

Hence

$$(4.5) \quad G_n^*(0) \rightarrow \frac{1}{k+1}.$$

We claim that $G(\zeta)$ is a meromorphic function on $\mathbb{C} \setminus E_2$. Suppose that $G(\zeta) \equiv \infty$. Obviously, $G_n^*(\zeta)$ have no zeros in Δ . Applying the maximum principle to the sequence $\frac{1}{G_n^*(\zeta)}$ of analytic functions, we see that $G_n^*(\zeta) \equiv \infty$ in Δ which contradict $G_n^*(0) \Rightarrow \frac{1}{k+1}$.

We claim that $G(\zeta) = \frac{\zeta^{k+1}}{k+1}$ on $\mathbb{C} \setminus E_2$. Indeed, since $G(\zeta)$ is a meromorphic function in $\mathbb{C} \setminus E_2$, then by Lemma 3.5, $G(\zeta) = \frac{\zeta^{k+1+c}}{k+1}$, where c is a constant. Since $G_n^*(\zeta)$ have no zeros in Δ , we have

$$G_n^*(\zeta) \Rightarrow \frac{\zeta^{k+1+c}}{k+1} \frac{1}{\zeta^{k+1}}, \quad \zeta \in \Delta.$$

Hence, $G_n^*(0) \rightarrow \frac{\zeta^{k+1+c}}{k+1} \frac{1}{\zeta^{k+1}} \Big|_{\zeta=0}$. By (4.5), we get that $c = 0$.

Suppose that $1 \notin E_2$. Since $G_n(1) = 0$, we have $G(1) = 0$ which contradicts that $G(\zeta) = \frac{\zeta^{k+1}}{k+1}$. Thus, $1 \in E_2$. By Lemma 3.8, $G(\zeta) = \int_1^\zeta \xi^k d\xi = \frac{\zeta^{k+1}-1}{k+1}$ which contradicts that $G(\zeta) = \frac{\zeta^{k+1}}{k+1}$.

Case 2.2: $1 \in E_1$

By Lemma 3.8,

$$(4.6) \quad F(\zeta) = \int_1^\zeta \xi^k d\xi = \frac{\zeta^{k+1}-1}{k+1}, \quad \zeta \in \mathbb{C} \setminus E_1.$$

Let e_j be the j th root of the equation $\zeta^{k+1}-1=0$, $j=1, 2, \dots, k+1$.

Claim. $E_1 = \{e_1, e_2, \dots, e_{k+1}\}$.

Proof. Suppose that $\zeta_0 \notin E_1$, where $\zeta_0^{k+1}-1=0$. Obviously, ζ_0 is a zero of $F(\zeta)$ but not a multiple zero of $F(\zeta)$ which contradicts that all of zeros of $F(\zeta)$ are multiple.

Suppose that $\zeta_0 \in E_1$, where $\zeta_0^{k+1}-1 \neq 0$. By Lemma 3.8, $F(\zeta) = \int_{\zeta_0}^\zeta \xi^{k+1} d\xi = \frac{\zeta^{k+1}-\zeta_0^{k+1}}{k+1}$, $\zeta \in \mathbb{C} \setminus E_1$. By (4.6), $\zeta_0^{k+1}=1$, a contradiction. $\square$

By Lemma 3.8, there exists $\delta_j > 0$ such that for sufficiently large n , $F_n(\zeta)$ have a single zero $\zeta_{n,j} \rightarrow e_j$ of order 2 and a single pole $\eta_{n,j} \rightarrow e_j$ of order 1 in $\Delta(e_j, \delta_j)$.

Set $z_{n,j} = a_n \zeta_{n,j}$. Thus, $f_n(z_{n,j}) = 0$ and $z_{n,j} \rightarrow 0$ as $n \rightarrow \infty$, where $j = 1, 2, \dots, k+1$. Set $B_n = \{z_{n,1}, z_{n,2}, \dots, z_{n,k+1}\}$. Do as in Case 2.1, we may assume that $b_n \rightarrow 0$ is the zero of f_n of smallest modulus in $\Delta \setminus B_n$. Set $r_n = \frac{a_n}{b_n}$. Obviously, $F_n(\frac{1}{r_n}) = 0$. Since $b_n \notin B_n$, $\frac{1}{r_n} \neq \zeta_{n,j}$, where $j = 1, 2, \dots, k+1$. Since $F_n(\zeta)$ have a single zero $\zeta_{n,j} \rightarrow e_j$ of order 2 in $\Delta(e_j, \delta_j)$, by Hurwitz's Theorem and (4.6) we have $\frac{1}{r_n} \rightarrow \infty$ and hence $r_n \rightarrow 0$ as $n \rightarrow \infty$.

Let $G_n(\zeta) = \frac{f_n(b_n \zeta)}{b_n^{k+1}}$. We have that for sufficiently large n ,

- (f1) $G_n(\zeta)$ have only $k+1$ zeros $r_n \zeta_{n,i}$ of order 2 and at least $k+1$ poles $r_n \eta_{n,i}$ of order 1 in Δ . Obviously, $|r_n \zeta_{n,i}| \rightarrow 0$ and $|r_n \eta_{n,i}| \rightarrow 0$ as $n \rightarrow \infty$;
- (f2) all zeros of $G_n(\zeta)$ are multiple;
- (f3) $G'_n(\zeta) \neq \zeta^k \widehat{h}(b_n \zeta)$ and $G_n(1) = 0$.

By Lemma 2.7, $\{G_n\}$ is normal in Δ' . By Lemma 3.9, $\{G_n\}$ is quasinormal in $\mathbb{C}$. Thus, there exist a subsequence of $\{G_n(\zeta)\}$ (that we continue to call $\{G_n(\zeta)\}$) and $E_3 \subset \mathbb{C}$ such that

- (g1) E_3 have no accumulation point in $\mathbb{C}$;
- (g2) $G_n(\zeta) \xrightarrow{X} G(\zeta)$ on $\mathbb{C} \setminus E_3$;
- (g3) for each $\zeta_0 \in E_3$, no subsequence of $\{G_n(\zeta)\}$ is normal at ζ_0 .

Obviously, $E_3 \cap \Delta' = \emptyset$ and all zeros of $G(\zeta)$ are multiple in $\mathbb{C} \setminus E_3$;

Let

$$G_n^*(\zeta) = G_n(\zeta) \frac{\prod_{j=1}^{k+1} (\zeta - r_n \eta_{n,j})}{\prod_{j=1}^{k+1} (\zeta - r_n \zeta_{n,j})^2},$$

$$F_n^*(\zeta) = F_n(\zeta) \frac{\prod_{j=1}^{k+1} (\zeta - \eta_{n,j})}{\prod_{j=1}^{k+1} (\zeta - \zeta_{n,j})^2}.$$

By (4.6),

$$G_n^*(r_n \zeta) = F_n^*(\zeta) \Rightarrow \frac{1}{k+1}, \quad \zeta \in \mathbb{C}.$$

Hence

$$(4.7) \quad G_n^*(0) \rightarrow \frac{1}{k+1}.$$

We claim that $G(\zeta)$ is a meromorphic function on $\mathbb{C} \setminus E_3$. Suppose that $G(\zeta) \equiv \infty$. Obviously, $G_n^*(\zeta)$ have no zeros in Δ . Applying the maximum principle to the sequence $\frac{1}{G_n^*(\zeta)}$ of analytic functions, we see that $G_n^*(\zeta) \equiv \infty$ in Δ which contradict $G_n^*(0) \Rightarrow \frac{1}{k+1}$.

We claim that $G(\zeta) = \frac{\zeta^{k+1}}{k+1}$ on $\mathbb{C} \setminus E_3$. Since $G(\zeta)$ is a meromorphic function in $\mathbb{C} \setminus E_3$, by Lemma 3.5, $G(\zeta) = \frac{\zeta^{k+1} + c}{k+1}$, where c is a constant. Since $G_n^*(\zeta)$ have no zeros in Δ , we have

$$G_n^*(\zeta) \Rightarrow \frac{\zeta^{k+1} + c}{k+1} \frac{1}{\zeta^{k+1}}, \quad \zeta \in \Delta$$

If $c \neq 0$ then we get $G_n^*(0) \rightarrow \infty$, and this contradicts (4.7).

We claim that $1 \in E_3$. Indeed, suppose that $1 \notin E_3$. Since $G_n(1) = 0$, we have $G(1) = 0$ which contradicts that $G(\zeta) = \frac{\zeta^{k+1}}{k+1}$. Thus, $1 \in E_3$. By Lemma 3.8, $G(\zeta) = \int_1^\zeta \zeta^k d\zeta = \frac{\zeta^{k+1} - 1}{k+1}$ which contradicts that $G(\zeta) = \frac{\zeta^{k+1}}{k+1}$.

Case 3: $h(0) = \infty$.

Suppose that 0 is a pole of order k of $h(z)$. Let us assume, making standard normalizations, that for $z \in \Delta$

$$h(z) = \frac{1}{z^k} + \frac{a_{k-1}}{z^{k-1}} + \cdots = \frac{\widehat{h}(z)}{z^k},$$

where $\widehat{h}(z) \neq 0, \infty$ in Δ and $\widehat{h}(0) = 1$.

Case 3.1: $k = 1$.

By Lemma 2.5 (with $\alpha = 0$), there exists points $z_n \rightarrow 0$, positive numbers $\rho_n \rightarrow 0$ and a subsequence of $\{f_n\}$ (that we continue to call $\{f_n\}$) such that

$$(4.8) \quad g_n(\zeta) = f_n(z_n + \rho_n \zeta) \xrightarrow{\chi} g(\zeta), \quad \zeta \in \mathbb{C},$$

where $g(\zeta)$ is a nonconstant meromorphic function in $\mathbb{C}$, all of whose zeros are multiple.

Again we consider two cases.

Case 1: $z_n/\rho_n \rightarrow \infty$.

Consider

$$(4.9) \quad \varphi_n(\zeta) = f_n(z_n + z_n \zeta) = f_n(z_n(1 + \zeta)).$$

Since g is nonconstant, there exist $\zeta_1, \zeta_2 \in \mathbb{C}$ such that $g(\zeta_1) \neq g(\zeta_2)$. We have for $j = 1, 2$,

$$g(\zeta_j) = \lim_{n \rightarrow \infty} f_n(z_n + \rho_n \zeta_j) = \lim_{n \rightarrow \infty} f_n(z_n + z_n(\frac{\rho_n}{z_n} \zeta_j)) = \lim_{n \rightarrow \infty} \varphi_n(\frac{\rho_n}{z_n} \zeta_j).$$

Since $(\frac{\rho_n}{z_n})\zeta_j \rightarrow 0$ as $n \rightarrow \infty$, the family $\{\varphi_n\}$ is not equicontinuous at 0 and hence no subsequence of $\{\varphi_n\}$ is normal at 0. By (4.9),

$$(4.10) \quad \varphi_n'(\zeta) = z_n f_n'(z_n(1 + \zeta)) \neq \frac{\widehat{h}(z_n(1 + \zeta))}{1 + \zeta}.$$

Since all zeros of φ_n are multiple and by (4.10), the family $\{\varphi_n\}$ is quasinormal in $\mathbb{C}$ by Lemma 3.9. Hence there exist a set E and a subsequence of $\{\varphi_n\}$ (that we continue to denote by $\{\varphi_n\}$) such that

(h1) E has no accumulation in $\mathbb{C}$;

(h2) $\varphi_n(\zeta) \xrightarrow{\chi} \varphi(\zeta)$ in $\mathbb{C} \setminus E$ and all the zeros of $\varphi_n(\zeta)$ are multiple;

(h3) for each $\zeta_0 \in E$, no subsequence of $\{\varphi_n\}$ is normal at ζ_0 .

Obviously, $0 \in E$. by Lemma 3.8, $\varphi(\zeta) = \int_0^\zeta \frac{1}{1+\xi} d\xi$. Hence, $\int_0^\zeta \frac{1}{1+\xi} d\xi$ is a multi-valued function in $\mathbb{C} \setminus E$. A contradiction.

Case 2: $\frac{z_n}{\rho_n} \rightarrow \alpha \in \mathbb{C}$.

Since $g'_n(\zeta) \neq \frac{\rho_n \hat{h}(z_n + \rho_n \zeta)}{z_n + \rho_n \zeta}$ and $\frac{\rho_n \hat{h}(z_n + \rho_n \zeta)}{z_n + \rho_n \zeta} \xrightarrow{\chi} \frac{1}{\zeta + \alpha}$ in $\mathbb{C}$, then by Lemma 3.4, either $g'(\zeta) \equiv \frac{1}{\alpha + \zeta}$ in $\mathbb{C}$, or $g'(\zeta) \neq \frac{1}{\alpha + \zeta}$ in $\mathbb{C}$. However, since all poles of g' are multiple, the first alternative obviously cannot hold, Thus $g'(\zeta) \neq \frac{1}{\alpha + \zeta}$. By Theorem E we deduce that g is rational. By the fundamental theorem of algebra, g cannot be a polynomial. hence

$$g'(\zeta) = \frac{1}{a + \zeta} + \frac{1}{p(\zeta)},$$

where $p(\zeta)$ is a polynomial. It is easy to see that if $\deg p \leq 1$, then the right hand side above cannot be a derivative of a nonconstant meromorphic function in $\mathbb{C}$. But then

$$g'(\zeta) = \frac{1}{\zeta} + O\left(\frac{1}{|\zeta|^2}\right), \quad \zeta \rightarrow \infty,$$

so that

$$\frac{1}{2\pi i} \int_{\Gamma(0, R)} g'(\zeta) d\zeta = 1 + o(R), \quad R \rightarrow \infty$$

for R sufficiently large, where $\Gamma(0, R)$ is the positive oriented circle of radius R about the origin. On the other hand, the above integral must vanish, as it is the integral of a derivative over a closed curve. A contradiction.

Case 3.2: $k \geq 2$.

We claim that for each $\delta > 0$, there exists at least one pole of f_n in $\Delta(0, \delta)$ for sufficiently large n . Otherwise, there exists a subsequence of $\{f_n\}$ (that we continue to call $\{f_n\}$) such that $\{f_n(z)\}$ is a sequence of functions holomorphic in $\Delta(0, \delta)$. By Lemma 3.11, $\{f_n\}$ is normal at 0. A contradiction.

Hence, taking a subsequence and renumbering if necessary, we may assume that $a_n \rightarrow 0$ is the pole of $f_n(z)$ of smallest modulus. Since $f'_n(z) \neq h(z)$ and $h(0) = \infty$, we have $f(0) \neq \infty$ and hence $a_n \neq 0$. Let $F_n(\zeta) = a_n^{k-1} f_n(a_n \zeta)$, we have

- (i1) $F_n(\zeta)$ is holomorphic function in Δ ;
- (i2) all zeros of $F_n(\zeta)$ are multiple;
- (i3) $F'_n(\zeta) \neq \frac{\hat{h}(a_n \zeta)}{\zeta^k}$ and $F_n(1) = \infty$.

By Lemma 3.11, $\{F_n(\zeta)\}$ is normal at Δ . By Lemma 3.9, $\{F_n(\zeta)\}$ is quasinormal in $\mathbb{C}$. Thus, there exists a subsequence of $\{F_n(\zeta)\}$ (that we continue to call $\{F_n(\zeta)\}$) and $E_4 \subset \mathbb{C}$ such that

- (j1) E have no accumulation point in $\mathbb{C}$;
- (j2) $F_n(\zeta) \xrightarrow{\chi} F(\zeta)$ in $\mathbb{C} \setminus E_4$;
- (j3) for each $\zeta_0 \in E_4$, no subsequence of $\{F_n(\zeta)\}$ is normal at ζ_0 .

Obviously, $E_4 \cap \Delta = \emptyset$ and all zeros of $F(\zeta)$ are multiple in $\mathbb{C} \setminus E_4$.

Claim. For n sufficiently large, $F_n(0) \neq 0$ (and hence $f_n(0) \neq 0$).

Proof. Otherwise, there exists a subsequence of $\{F_n(\zeta)\}$ (that we continue to call $\{F_n(\zeta)\}$) such that $F_n(0) = 0$. Thus $F(0) = 0$ and hence $F(\zeta)$ is a meromorphic function in $\mathbb{C} \setminus E_4$. Suppose that E_4 is the empty set. Since $F'_n(\zeta) \neq \frac{\hat{h}(a_n\zeta)}{\zeta^k}$ and $\frac{\hat{h}(a_n\zeta)}{\zeta^k} \xrightarrow{\chi} \frac{1}{\zeta^k}$ in $\mathbb{C}$, by Lemma 3.4 we have either $F'(\zeta) \equiv \frac{1}{\zeta^k}$ in $\mathbb{C}$, or $F'(\zeta) \neq \frac{1}{\zeta^k}$ in $\mathbb{C}$. If $F'(\zeta) - \frac{1}{\zeta^k} \equiv 0$ in $\mathbb{C}$, then we have $F(0) = \infty$ which contradicts that $F(0) = 0$. If $F'(\zeta) \neq \frac{1}{\zeta^k}$ in $\mathbb{C}$, then, by Theorem E and Lemma 3.1, we have $F(\zeta) = c$, where $c \in \mathbb{C}$ is a constant. On the other hand, since $F_n(1) = \infty$, we have $F(1) = \infty$, a contradiction.

Thus, we can assume that E_4 is not the empty set. Suppose that $\zeta_0 \in E_4$. Obviously, $\zeta_0 \neq 0$. By Lemma 3.8, we have $F(\zeta) = \int_{\zeta_0}^{\zeta} \frac{1}{\xi^k} d\xi$ which implies $F(0) = \infty$, a contradiction to the fact that $F_n(0) = 0$. $\square$

Without loss of generality, we assume that for each $n \in \mathbb{N}$, $f_n(0) \neq 0$.

We claim that for each $\delta > 0$, there exists at least one zero of f_n in $\Delta(0, \delta)$ for sufficiently large n . Otherwise, there exists a subsequence of $\{f_n\}$ (that we continue to call $\{f_n\}$) such that $f_n(z) \neq 0$ in $\Delta(0, \delta)$. Since $f_n(z) \neq h(z)$, we have f_n and h have no common poles for each n and $f_n(z) - h(z) \neq 0$. By Lemma 2.3, $\{f_n\}$ is normal at 0. A contradiction.

Hence, taking a subsequence and renumbering if necessary, we may assume that b_n be the zero of $\{f_n\}$ of smallest modulus. Obviously we have $b_n \rightarrow 0$ as $n \rightarrow \infty$. Since $f_n(0) \neq 0$, we have $b_n \neq 0$. Let $G_n(\zeta) = b_n^{k-1} f_n(b_n \zeta)$, we have

- (k1) $G_n(\zeta) \neq 0$ in Δ ;
- (k2) all zeros of $G_n(\zeta)$ are multiple;
- (k3) $G'_n(\zeta) \neq \frac{\hat{h}(b_n \zeta)}{\zeta^k}$ and $G_n(1) = 0$.

By Lemma 3.10, $\{G_n(\zeta)\}$ is normal at Δ . By Lemma 3.9, $\{G_n(\zeta)\}$ is quasinormal in $\mathbb{C}$. Thus, there exists a subsequence of $\{G_n(\zeta)\}$ (that we continue to call $\{G_n(\zeta)\}$) and $E_5 \subset \mathbb{C}$ such that

- (l1) E_5 have no accumulation point in $\mathbb{C}$;
- (l2) $G_n(\zeta) \xrightarrow{\chi} G(\zeta)$ on $\mathbb{C} \setminus E_5$;
- (l3) for each $\zeta_0 \in E_5$, no subsequence of $\{G_n(\zeta)\}$ is normal at ζ_0 .

Obviously, $E_5 \cap \Delta = \emptyset$ and all zeros of $G(\zeta)$ are multiple in $\mathbb{C} \setminus E_5$.

Case 3.2.1: E_5 is the empty set.

Since $G_n(1) = 0$, then $G(1) = 0$ and hence 1 is a multiple zero of $G(\zeta)$ and $G(\zeta)$ is a meromorphic function. Since $G'_n(\zeta) \neq \frac{\hat{h}(b_n \zeta)}{\zeta^k}$ and $\frac{\hat{h}(b_n \zeta)}{\zeta^k} \xrightarrow{\chi} \frac{1}{\zeta^k}$ in $\mathbb{C}$, then by Lemma 3.4, we have either $G'(\zeta) \equiv \frac{1}{\zeta^k}$ in $\mathbb{C}$, or $G'(\zeta) \neq \frac{1}{\zeta^k}$ in $\mathbb{C}$.

If $G'(\zeta) \equiv \frac{1}{\zeta^k}$ in $\mathbb{C}$, then we have $G(\zeta) = \frac{1}{1-k} (\frac{1}{\zeta^{k-1}} + c)$. Obviously, 1 is not a multiple zero of $G(\zeta)$, a contradiction.

If $G'(\zeta) \neq \frac{1}{\zeta^k}$ in $\mathbb{C}$, then by Theorem E and Lemma 3.1, we have $G(\zeta) = c$. Since $G(1) = 0$, we have $G(\zeta) \equiv 0$. i.e.,

$$(4.11) \quad G_n(\zeta) = b_n^{k-1} f_n(b_n \zeta) \Rightarrow 0, \quad \text{in } \mathbb{C}.$$

Since $G_n(\frac{a_n}{b_n}) = b_n^{k-1}f_n(a_n) = \infty$, by Hurwitz's Theorem and (4.11), we have $\frac{a_n}{b_n} \rightarrow \infty$. Writing $r_n = \frac{b_n}{a_n}$. Hence $r_n \rightarrow 0$ as $n \rightarrow \infty$.

Since $F_n(r_n) = a_n^{k-1}f_n(b_n) = 0$, we have $F(0) = 0$ and hence $F(z)$ is a meromorphic function in $\mathbb{C} \setminus E_4$. Since $F'_n(\zeta) \neq \frac{\widehat{h}(a_n\zeta)}{\zeta^k}$ and $\frac{\widehat{h}(a_n\zeta)}{\zeta^k} \xrightarrow{\chi} \frac{1}{\zeta^k}$ in $\mathbb{C}$, we have by Lemma 3.4 either $F'(\zeta) \equiv \frac{1}{\zeta^k}$ in $\mathbb{C} \setminus E_4$, or $F'(\zeta) \neq \frac{1}{\zeta^k}$ in $\mathbb{C} \setminus E_4$.

Now, we consider the following two cases.

Case 1: E_4 is the empty set.

If $F'(\zeta) - \frac{1}{\zeta^k} \equiv 0$ in $\mathbb{C}$, then we have $F(0) = \infty$ which contradicts that $F(0) = 0$. If $F'(\zeta) \neq \frac{1}{\zeta^k}$ in $\mathbb{C}$, then by Theorem E and Lemma 3.1, we have $F(\zeta) = c$ which contradicts that $F(1) = \infty$.

Case 2: E_4 is not the empty set.

Let $\zeta_0 \in E$. Obviously, $\zeta_0 \neq 0$. By Lemma 3.8, we have $F(\zeta) = \int_{\zeta_0}^{\zeta} \frac{1}{\xi^k} d\xi = \frac{1}{1-k} \left(\frac{1}{\zeta^{k-1}} - \frac{1}{\zeta_0^{k-1}} \right)$. Obviously, we have $F(0) = \infty$ which contradicts $F(0) = 0$.

Case 3.2.2: E_5 is not the empty set.

Let $\zeta_0 \in E_5$. Obviously, $\zeta_0 \neq 0$. By Lemma 3.8, we have

$$G(\zeta) = \int_{\zeta_0}^{\zeta} \frac{1}{\xi^k} d\xi = \frac{1}{(k-1)\zeta_0^{k-1}} \left(\frac{\zeta^{k-1} - \zeta_0^{k-1}}{\zeta^{k-1}} \right), \quad \zeta \in \mathbb{C} \setminus E_5.$$

Let ζ_i ($i = 1, 2, \dots, k-1$) be the $k-1$ roots of the equation $\zeta^{k-1} - \zeta_0^{k-1} = 0$.

Do as in Case 2.2, we have $E_5 = \{\zeta_1, \zeta_2, \dots, \zeta_{k-1}\}$.

We claim that $1 \in E_5$ and hence $G(\zeta) = \frac{1}{1-k} \left(\frac{1}{\zeta^{k-1}} - 1 \right)$. Indeed, if $1 \notin E_5$, i.e., $1 - \zeta_0^{k-1} \neq 0$, then $G(1) = \frac{\zeta_0^{k-1} - 1}{(1-k)\zeta_0^{k-1}} \neq 0$. On the other hand, since $G_n(1) = 0$, we have $G(1) = 0$, a contradiction.

Now we have

$$(4.12) \quad G_n(\zeta) \xrightarrow{\chi} \frac{\zeta^{k-1} - 1}{(k-1)\zeta^{k-1}}, \quad \zeta \in \Delta.$$

By Hurwitz's Theorem, there exist $\gamma_{n,i}$, $i = 1, 2, \dots, k-1$ such that $\gamma_{n,i} \rightarrow 0$ and $G_n(\gamma_{n,i}) = \infty$. Since f and h have no common poles, then $G_n(0) \neq \infty$, and we have that $\gamma_{n,i} \neq 0$ ($i = 1, 2, \dots, k-1$).

Let s_n be one of $\{\gamma_{n,1}, \gamma_{n,2}, \dots, \gamma_{n,k-1}\}$ of largest modulus. Set $U_n(\xi) = s_n^{k-1}G_n(s_n\xi)$. Obviously, for n sufficiently large, $U_n(\xi)$ have only $k-1$ poles $\eta_{n,i} = \frac{\gamma_{n,i}}{s_n}$ on $\bar{\Delta}$. By (4.12), we have for each $R > 0$,

$$U_n(\xi) \neq 0 \quad \xi \in \Delta(0, R)$$

for n sufficiently large. By (k_3) $U'_n(\xi) \neq \frac{\widehat{h}(b_n s_n \xi)}{\xi^k}$, and then we get by Lemma 3.10 that $U_n(\xi)$ is normal in $\mathbb{C}$. Without loss of generality, we assume that $U_n(\xi) \xrightarrow{\chi} U(\xi)$ in $\mathbb{C}$ and $\eta_{n,i} \rightarrow \eta_i$. Since $U_n(1) = \infty$, we have $U(1) = \infty$.

We claim that $U(\xi) \equiv \infty$ in $\mathbb{C}$. Otherwise, by Lemma 3.4, either $U'(\xi) \equiv \frac{1}{\xi^k}$ in $\mathbb{C}$, or $U'(\xi) \neq \frac{1}{\xi^k}$ in $\mathbb{C}$. If $U'(\xi) \equiv \frac{1}{\xi^k}$ in $\mathbb{C}$, then we have $U(\xi) = \frac{1}{1-k} \left(\frac{1}{\xi^{k-1}} + c \right)$ which

contradicts that $U(1) = \infty$. If $U'(\xi) \neq \frac{1}{\xi^k}$ on $\mathbb{C}$, then, by Theorem E and Lemma 3.1, we have $U(\xi) = c$ which contradicts $U(1) = \infty$.

Now, we have

$$U_n(\xi) \xrightarrow{\chi} \infty, \quad \xi \in \mathbb{C}.$$

Let $U_n^*(\xi) = U_n(\xi) \cdot \prod_{i=1}^{k-1} (\xi - \eta_{n,i})$. By the maximum principle applied to U_n^* , we get that

$$U_n^*(\xi) \xrightarrow{\chi} \infty, \quad \xi \in \mathbb{C}.$$

Let $G_n^*(\zeta) = G_n(\zeta) \cdot \prod_{i=1}^{k-1} (\zeta - \gamma_{n,i}) = G_n(\zeta) \cdot \prod_{i=1}^{k-1} (\zeta - s_n \eta_{n,i})$. By (4.12), for sufficiently large n , $G_n^*(\zeta)$ have no pole in $\Delta(0, \frac{1}{2})$ and hence, by the maximum principle,

$$G_n^*(\zeta) \xrightarrow{\chi} \frac{\zeta^{k-1} - 1}{k - 1}, \quad \zeta \in \Delta(0, \frac{1}{2}).$$

So we have $G_n^*(0) \rightarrow \frac{1}{1-k}$ as $n \rightarrow \infty$. On the other hand,

$$\begin{aligned} G_n^*(s_n \xi) &= G_n(s_n \xi) \cdot \prod_{i=1}^{k-1} (s_n \xi - \gamma_{n,i}) = G_n(s_n \xi) \cdot \prod_{i=1}^{k-1} (s_n \xi - s_n \eta_{n,i}) \\ (4.13) \quad &= s_n^{k-1} G_n(s_n \xi) \cdot \prod_{i=1}^{k-1} (\xi - \eta_{n,i}) = U_n^*(\xi) \xrightarrow{\chi} \infty, \quad \xi \in \mathbb{C}. \end{aligned}$$

(4.13) implies that $G_n^*(0) \rightarrow \infty$ as $n \rightarrow \infty$ which contradicts $G_n^*(0) \rightarrow \frac{1}{1-k}$ as $n \rightarrow \infty$. The proof of the Theorem is complete. $\square$

REFERENCES

- [1] Naum Il'yich Akhiezer, *Elements of the Theory of Elliptic Functions*, (1970) Moscow, translated into English as AMS Translations of Mathematical Monographs Volume 79 (1990) AMS, Rhode Island ISBN 0-8218-4532-2.
- [2] Steven B. Bank and J. K. Langley, *On the value distribution theory of elliptic functions*, Monatshefte der Math. **98**, no. 1 (1984), 1-20.
- [3] W. Bergweiler and A. Eremenko, *On the singularities of the inverse to a meromorphic function of finite order*, Rev. Mat. Iberoamericana **11** (1995), 355-373.
- [4] W. Bergweiler and Xuecheng Pang, *On the derivative of meromorphic functions with multiple zeros*, J. Math. Anal. Appl. **278** (2003), 285-292.
- [5] J.M. Chang and L. Zalcman, *Meromorphic functions that share a set with their derivatives*, J. Math. Anal. Appl. **338** (2008), 1020-1028.
- [6] John B. Conway, *Functions of One Complex Variable*, Springer-Verlag, New York, 1973.
- [7] W. K. Hayman, *Picard values of meromorphic functions and their derivatives*, Ann. Math. (2) **70** (1959), 9-42.
- [8] X.J. Liu and S. Nevo, *A criterion of normality established by a single holomorphic function*, Acta Math. Sin., English Series, to appear.
- [9] X.J. Liu, S. Nevo and X.C. Pang, *On the k -th derivative of meromorphic functions with zeros of multiplicity at least $k + 1$* , J. Math. Anal. Appl. **348** (2008), 516-529.
- [10] S. Nevo, *Applications of Zalcman's Lemma to Q_m -normal families*, Analysis (Munich) **21** (2001), 289-325.

- [11] Shahar Nevo, On Theorems of Yang and Schwick, *Complex Variables and Elliptic Equations* **46**, no. 4 (2001), 315-321.
- [12] S. Nevo, *Applications of Zalcman's Lemma to Q_m -normal families*, Analysis (Munich) **21** (2001), 289-325.
- [13] Shahar Nevo, Xuecheng Pang and Lawrence Zalcman, *Picard-Hayman behavior of derivatives of meromorphic functions with multiple zeros*, Electron. Res. Announce. Amer. Math. Soc. **12** (2006), 37-43.
- [14] Shahar Nevo, Xuecheng Pang and Lawrence Zalcman, *Quasinormality and meromorphic functions with multiple zeros*, J. D'Anal. Math. **101** (2007), 1-23.
- [15] Xuecheng Pang and Lawrence Zalcman, *Normal families and shared values*, Bull. London Math. Soc. **32** (2000), 325-331.
- [16] Xuecheng Pang, Degui Yang and Lawrence Zalcman, *Normal families and omitted functions*, Indiana Univ. Math. J. **54** (2005), 223-235.
- [17] Xuecheng Pang, Shahar Nevo and Lawrence Zalcman, *Quasinormal families of meromorphic functions II*, Operator Theory: Advances and Applications **158** (2005), 177-189.
- [18] Xuecheng Pang, Mingliang Fang, and Lawrence Zalcman, *Normal families of holomorphic functions with multiple zeros*, Conform. Geom. Dyn. **11** (2007), 101-106.
- [19] Xuecheng Pang, Shahar Nevo and Lawrence Zalcman, *Derivatives of meromorphic functions with multiple zeros and rational functions*, Comput. Methods Funct. Theory **8**, no.2 (2008), 483-491.
- [20] W. Schwick, *On Hayman's alternative for families of meromorphic functions*, Complex. Variables **32** (1997), 51-57.
- [21] M. Tsuji, *On Borel's directions of meromorphic functions of finite order. II*, Kodai Math. Sem. Rep. **2**, Nos. 4-5 (1950), 96-100.
- [22] M. Tsuji, *Potential Theory in Modern Function Theory*, 2nd Ed. Chelsea, 1975.
- [23] Yuefei Wang and Mingliang Fang, *Picard values and normal families of meromorphic functions with multiple zeros*, Acta Math. Sinica, New Ser. **14** (1998), 17-26.

E-mail address: yangpai@cuit.edu.cn

DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, 52900 RAMAT-GAN, ISRAEL

E-mail address: nevosh@macs.biu.ac.il

DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200062, P. R. CHINA

E-mail address: xcpang@euler.math.ecnu.edu.cn